

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



# (12) UK Patent Application (19) GB (11) 2 302 689 (13) A

(43) Date of A Publication 29.01.1997

(21) Application No 9613003.4

(22) Date of Filing 20.06.1996

(30) Priority Data

(31) 9513118

(32) 28.06.1995

(33) GB

(71) Applicant(s)

Merck Sharp & Dohme Limited

(Incorporated in the United Kingdom)

Hertford Road, HODDESDON, Hertfordshire,  
EN11 9BU, United Kingdom

(72) Inventor(s)

Jonathan David Moseley

Christopher John Swain

Brian John Williams

(74) Agent and/or Address for Service

I J Hiscock

Merck & Co. Inc, European Patent Department,  
Terlings Park, Eastwick Road, HARLOW, Essex,  
CM20 2QR, United Kingdom

(51) INT CL<sup>6</sup>

C07D 413/06, A61K 31/535, C07D 413/14

(52) UK CL (Edition O)

C2C CAA CLZ CQT C1341 C1450 C1562 C215 C22Y  
C220 C248 C25Y C250 C251 C252 C255 C28X C31Y  
C311 C32Y C323 C334 C36Y C364 C37Y C373 C464  
C613 C620 C650 C672 C694 C697 C699 C80Y C802  
U1S S1321 S1328 S2414 S2415 S2416 S2417

(56) Documents Cited

None

(58) Field of Search

UK CL (Edition O) C2C  
INT CL<sup>6</sup> C07D  
ONLINE: CAS

## (54) N-oxides of morpholine derivatives and their use as therapeutic agents


(57) The present invention relates to morpholine derivatives of the formula (I) and pharmaceutically acceptable salts thereof wherein

R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup>, R<sup>9a</sup>, R<sup>9b</sup>, X, Y, Z and Het are as defined in the specification;

m is 0 or 1; and

n is 0 or 1, where the sum total of n+m is 1 or 2.

The compounds are of particular use in the treatment of pain, inflammation, migraine, emesis and postherpetic neuralgia.



(I)

32  
32  
64

GB 2 302 689 A

N-OXIDES OF MORPHOLINE DERIVATIVES AND THEIR USE AS  
THERAPEUTIC AGENTS

5 This invention relates to a class of morpholine derivatives which are useful as tachykinin antagonists. More particularly, this invention relates to the N-oxides of a class of morpholine derivatives which contain an amine-substituted azo-heterocyclic moiety.

10 The tachykinins are a group of naturally occurring peptides found widely distributed throughout mammalian tissues, both within the central nervous system and in peripheral nervous and circulatory systems.

The tachykinins are distinguished by a conserved carboxyl-terminal sequence:

Phe-X-Gly-Leu-Met-NH<sub>2</sub>

15 At present, there are three known mammalian tachykinins referred to as substance P, neurokinin A (NKA, substance K, neuromedin L) and neurokinin B (NKB, neuromedin K) (for review see J.E. Maggio, *Peptides* (1985) 6(suppl. 3), 237-242). The current nomenclature designates the three tachykinin receptors mediating the biological actions of substance P, NKA and NKB as the NK<sub>1</sub>, NK<sub>2</sub> and NK<sub>3</sub> receptors, respectively.

20 Evidence for the usefulness of tachykinin receptor antagonists in pain, headache, especially migraine, Alzheimer's disease, multiple sclerosis, attenuation of morphine withdrawal, cardiovascular changes, oedema, such as oedema caused by thermal injury, chronic inflammatory diseases such as rheumatoid arthritis, asthma/bronchial hyperreactivity  
25 and other respiratory diseases including allergic rhinitis, inflammatory diseases of the gut including ulcerative colitis and Crohn's disease, ocular injury and ocular inflammatory diseases, proliferative vitreoretinopathy, irritable bowel syndrome and disorders of bladder function including cystitis and bladder detruser hyper-reflexia is reviewed in "Tachykinin  
30 Receptors and Tachykinin Receptor Antagonists", C.A. Maggi, R.

Patacchini, P. Rovero and A. Giachetti, *J. Auton. Pharmacol.* (1993) 13, 23-93.

For instance, substance P is believed *inter alia* to be involved in the neurotransmission of pain sensations [Otsuka *et al*, "Role of Substance P as a Sensory Transmitter in Spinal Cord and Sympathetic Ganglia" in 5 1982 Substance P in the Nervous System, *Ciba Foundation Symposium* 91, 13-34 (published by Pitman) and Otsuka and Yanagisawa, "Does Substance P Act as a Pain Transmitter?" *TIPS* (1987) 8, 506-510], specifically in the transmission of pain in migraine (B.E.B. Sandberg *et al*, 10 *J. Med Chem.*, (1982) 25, 1009) and in arthritis [Levine *et al* in *Science* (1984) 226, 547-549]. Tachykinins have also been implicated in gastrointestinal (GI) disorders and diseases of the GI tract such as inflammatory bowel disease [Mantyh *et al* in *Neuroscience* (1988) 25(3), 817-37 and D. Regoli in "Trends in Cluster Headache" Ed. Sicuteri *et al* 15 Elsevier Scientific Publishers, Amsterdam (1987) page 85)] and emesis [F. D. Tattersall *et al*, *Eur. J. Pharmacol.*, (1993) 250, R5-R6]. It is also hypothesised that there is a neurogenic mechanism for arthritis in which substance P may play a role [Kidd *et al* "A Neurogenic Mechanism for Symmetrical Arthritis" in *The Lancet*, 11 November 1989 and Grönblad *et al*, 20 "Neuropeptides in Synovium of Patients with Rheumatoid Arthritis and Osteoarthritis" in *J. Rheumatol.* (1988) 15(12), 1807-10]. Therefore, substance P is believed to be involved in the inflammatory response in diseases such as rheumatoid arthritis and osteoarthritis, and fibrositis [O'Byrne *et al*, *Arthritis and Rheumatism* (1990) 33, 1023-8]. Other 25 disease areas where tachykinin antagonists are believed to be useful are allergic conditions [Hamelet *et al*, *Can. J. Pharmacol. Physiol.* (1988) 66, 1361-7], immunoregulation [Lotz *et al*, *Science* (1988) 241, 1218-21 and Kimball *et al*, *J. Immunol.* (1988) 141(10), 3564-9] vasodilation, bronchospasm, reflex or neuronal control of the viscera [Mantyh *et al*, 30 *Proc. Natl. Acad. Sci., USA* (1988) 85, 3235-9] and, possibly by arresting or slowing  $\beta$ -amyloid-mediated neurodegenerative changes [Yankner *et al*,


*Science* (1990) 250, 279-82] in senile dementia of the Alzheimer type, Alzheimer's disease and Down's Syndrome.

Tachykinin antagonists may also be useful in the treatment of small cell carcinomas, in particular small cell lung cancer (SCLC) [Langdon *et al*, *Cancer Research* (1992) 52, 4554-7].

Substance P may also play a role in demyelinating diseases such as multiple sclerosis and amyotrophic lateral sclerosis [J. Lubner-Narod *et al*, poster *C.I.N.P. XVIIIth Congress*, 28th June-2nd July 1992], and in disorders of bladder function such as bladder detrusor hyper-reflexia (*The Lancet*, 16th May 1992, 1239).

It has furthermore been suggested that tachykinins have utility in the following disorders: depression, dysthymic disorders, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina and Reynauld's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, neuropathy, neuralgia, disorders related to immune enhancement or suppression such as systemic lupus erythmatosus (European patent specification no. 0 436 334), ophthalmic disease such as conjunctivitis, vernal conjunctivitis, and the like, and cutaneous diseases such as contact dermatitis, atopic dermatitis, urticaria, and other eczematooid dermatitis (European patent specification no. 0 394 989).

European patent specification no. 0 577 394 (published 5th January 1994) discloses morpholine and thiomorpholine tachykinin receptor antagonists of the general formula




wherein R<sup>1</sup> is a large variety of substituents;

R<sup>2</sup> and R<sup>3</sup> are inter alia hydrogen;

R<sup>4</sup> is inter alia

5



R<sup>5</sup> is inter alia optionally substituted phenyl;

R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are a variety of substituents;

10 X is O, S, SO or SO<sub>2</sub>;


Y is inter alia O; and

Z is hydrogen or C<sub>1-4</sub>alkyl.

We have now found a further class of non-peptides which are potent antagonists of tachykinins, especially of substance P.

15 It is desirable that compounds may be administered orally and by injection. Certain compounds have now been discovered which act as potent non-peptide tachykinin antagonists and which, by virtue of their advantageous aqueous solubility, are particularly easily formulated for administration by both the oral and injection routes, for example in  
20 aqueous media.

The present invention provides compounds of the formula (I):



wherein

$R^1$  is hydrogen, halogen,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkoxy,  $CF_3$ ,  $OCF_3$ ,  $NO_2$ , CN,  $SR^a$ ,  $SOR^a$ ,  $SO_2R^a$ ,  $CO_2R^a$ ,  $CONR^aR^b$ ,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl or  $C_{1-4}$ alkyl substituted by  $C_{1-4}$ alkoxy, wherein  $R^a$  and  $R^b$  each independently represent hydrogen or  $C_{1-4}$ alkyl;

$R^2$  is hydrogen, halogen,  $C_{1-6}$ alkyl,  $CF_3$ ,  $OCF_3$  or  $C_{1-6}$ alkoxy substituted by  $C_{1-4}$ alkoxy;

$R^3$  is hydrogen, halogen,  $CF_3$  or  $OCF_3$ ;

$R^4$  is hydrogen, halogen,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkoxy,  $CF_3$ ,  $NO_2$ , CN,  $SR^a$ ,  $SOR^a$ ,  $SO_2R^a$ ,  $CO_2R^a$ ,  $CONR^aR^b$ ,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl or  $C_{1-4}$ alkyl substituted by  $C_{1-4}$ alkoxy, wherein  $R^a$  and  $R^b$  are as previously defined;

$R^5$  is hydrogen, halogen,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkoxy substituted by  $C_{1-4}$ alkoxy or  $CF_3$ ;

$R^6$  is hydrogen,  $C_{1-4}$ alkyl,  $C_{3-7}$ cycloalkyl,  $C_{3-7}$ cycloalkyl $C_{1-4}$ alkyl, or  $C_{2-4}$ alkyl substituted by  $C_{1-4}$ alkoxy or hydroxy;

$R^7$  is hydrogen,  $C_{1-4}$ alkyl,  $C_{3-7}$ cycloalkyl,  $C_{3-7}$ cycloalkyl $C_{1-4}$ alkyl, or  $C_{2-4}$ alkyl substituted by one or two substituents selected from  $C_{1-4}$ alkoxy, hydroxy or a 4, 5 or 6 membered heteroaliphatic ring containing one or two heteroatoms selected from N, O and S;



or R<sup>6</sup> and R<sup>7</sup>, together with the nitrogen atom to which they are attached, form a saturated or partially saturated heterocyclic ring of 4 to 7 ring atoms, which ring may optionally contain in the ring one oxygen or sulphur atom or a group selected from NR<sup>8</sup>, S(O) or S(O)<sub>2</sub> and which ring  
5 may be optionally substituted by one or two groups selected from hydroxy, C<sub>1-4</sub>alkyl, hydroxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, oxo, COR<sup>a</sup> or CO<sub>2</sub>R<sup>a</sup> where R<sup>a</sup> is as previously defined;

or R<sup>6</sup> and R<sup>7</sup> together with the nitrogen atom to which they are attached, form a non-aromatic azabicyclic ring system of 6 to 12 ring  
10 atoms;

or Z, R<sup>6</sup> and the nitrogen atom to which they are attached form a heteroaliphatic ring of 4 to 7 ring atoms which may optionally contain an oxygen ring atom;

R<sup>8</sup> is hydrogen, C<sub>1-4</sub>alkyl, hydroxyC<sub>1-4</sub>alkyl or C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl;

15 R<sup>9a</sup> and R<sup>9b</sup> are each independently hydrogen or C<sub>1-4</sub>alkyl, or R<sup>9a</sup> and R<sup>9b</sup> are joined so, together with the carbon atoms to which they are attached, there is formed a C<sub>5-7</sub> ring;

Het is a 5- or 6-membered heterocyclic ring containing 2 or 3 nitrogen atoms optionally substituted by =O, =S or a C<sub>1-4</sub>alkyl group;

20 X is an alkylene chain of 1 to 4 carbon atoms optionally substituted by oxo;

Y is a C<sub>1-4</sub>alkyl group optionally substituted by hydroxy;

Z is C<sub>1-6</sub>alkylene or C<sub>3-7</sub>cycloalkylene;

m is 0 or 1; and

25 n is 0 or 1, where the sum total of n+m is 1 or 2;

and pharmaceutically acceptable salts thereof.

According to an alternative aspect of the present invention, R<sup>1</sup> is hydrogen, halogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, CF<sub>3</sub>, NO<sub>2</sub>, CN, SR<sup>a</sup>, SOR<sup>a</sup>, SO<sub>2</sub>R<sup>a</sup>, CO<sub>2</sub>R<sup>a</sup>, CONR<sup>a</sup>R<sup>b</sup>, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl or C<sub>1-4</sub>alkyl substituted by  
30 C<sub>1-4</sub>alkoxy, wherein R<sup>a</sup> and R<sup>b</sup> each independently represent hydrogen or

C<sub>1-4</sub>alkyl; R<sup>2</sup> is hydrogen, halogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy substituted by C<sub>1-4</sub>alkoxy or CF<sub>3</sub>; and R<sup>3</sup> is hydrogen, halogen or CF<sub>3</sub>.

A preferred class of compounds of formula (I) is that wherein R<sup>1</sup> is hydrogen, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, halogen or CF<sub>3</sub>.

5 Another preferred class of compounds of formula (I) is that wherein R<sup>2</sup> is hydrogen, C<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxy, halogen or CF<sub>3</sub>.

Also preferred is the class of compounds of formula (I) wherein R<sup>3</sup> is hydrogen, fluorine, chlorine or CF<sub>3</sub>.

10 A particularly preferred class of compounds of formula (I) is that wherein R<sup>1</sup> is fluorine, chlorine or CF<sub>3</sub>.

Another particularly preferred class of compounds of formula (I) is that wherein R<sup>2</sup> is hydrogen, fluorine, chlorine or CF<sub>3</sub>.

Also particularly preferred is the class of compounds of formula (I) wherein R<sup>3</sup> is hydrogen, fluorine, chlorine or CF<sub>3</sub>.

15 Preferably R<sup>1</sup> and R<sup>2</sup> are in the 3 and 5 positions of the phenyl ring.

More preferably R<sup>1</sup> is 3-fluoro or 3-CF<sub>3</sub>.

More preferably R<sup>2</sup> is 5-fluoro or 5-CF<sub>3</sub>.

More preferably R<sup>3</sup> is hydrogen.

Most preferably R<sup>1</sup> is 3-F or 3-CF<sub>3</sub>, R<sup>2</sup> is 5-CF<sub>3</sub> and R<sup>3</sup> is hydrogen.

20 A further preferred class of compound of formula (I) is that wherein R<sup>4</sup> is hydrogen.


Another preferred class of compounds of formula (I) is that wherein R<sup>5</sup> is hydrogen, fluorine, chlorine or CF<sub>3</sub>.

Preferably R<sup>4</sup> is hydrogen and R<sup>5</sup> is hydrogen or 4-fluoro.

25 Also preferred is the class of compounds of formula (I) wherein R<sup>9a</sup> and R<sup>9b</sup> are each independently hydrogen or methyl. Preferably R<sup>9a</sup> is hydrogen. Preferably R<sup>9b</sup> is hydrogen. Most preferably R<sup>9a</sup> and R<sup>9b</sup> are both hydrogen.

30 A further preferred class of compounds of formula (I) is that wherein m is 1.

From the foregoing it will be appreciated that a particularly apt sub-group of compounds of this invention are those of the formula (Ia) and pharmaceutically acceptable salts thereof:



5

wherein  $A^1$  is fluorine or  $CF_3$ ;

$A^2$  is fluorine or  $CF_3$ ;

$A^3$  is fluorine or hydrogen;

10 and  $R^6$ ,  $R^7$ , X, Y, Z, Het, m and n are as defined in relation to formula (I).

A preferred group Y for compounds of the formulae (I) or (Ia) is the  $C_{1-4}$ alkyl group, especially the methyl group.

Where the group  $NR^6R^7$  forms a saturated heterocyclic ring of 4 to 7 ring atoms which may optionally contain in the ring one oxygen or  
 15 sulphur atom or a group selected from  $NR^8$ ,  $S(O)$  or  $S(O)_2$ , suitable heterocyclic groups include azetidiny, pyrrolidino, piperidino, homopiperidino, piperazino, N-methylpiperazino, morpholino and thiomorpholino.

Suitable substituents on the saturated heterocyclic ring include  
 20  $CH_2OH$ ,  $CH_2OCH_3$ , oxo,  $CHO$ ,  $CO_2H$ ,  $CO_2CH_3$ , and  $CO_2CH_2CH_3$ .

When used herein the term "halogen" means fluorine, chlorine, bromine and iodine. The most apt halogen are fluorine and chlorine of which fluorine is preferred.

When used herein the term "alkyl" or "alkoxy" as a group or part of a group means that the group is straight or branched. Examples of suitable alkyl groups include methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl and t-butyl. Examples of suitable alkoxy groups include methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy and t-butoxy.

The term "alkenyl" as a group or part of a group means that the group is straight or branched and contains at least one double bond. Examples of suitable alkenyl groups include vinyl and allyl.

The term "alkynyl" as a group or part of a group means that the group is straight or branched and contains at least one triple bond. An example of a suitable alkynyl group is propargyl.

Suitable cycloalkyl and cycloalkyl-alkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl and cyclobutylmethyl.

Where the group  $NR^6R^7$  represents a heteroaliphatic ring of 4 to 7 ring atoms and said ring is partially saturated, a particularly preferred group is 3-pyrroline.

Where the group  $NR^6R^7$  represents a non-aromatic azabicyclic ring system, such a system may contain between 6 and 12, and preferably between 7 and 10, ring atoms. Suitable rings include 5-azabicyclo[2.1.1]hexyl, 5-azabicyclo[2.2.1]heptyl, 6-azabicyclo[3.2.1]octyl, 2-azabicyclo[2.2.2]octyl, 6-azabicyclo[3.2.2]nonyl, 6-azabicyclo[3.3.1]nonyl, 6-azabicyclo[3.2.2]decyl, 7-azabicyclo[4.3.1]decyl, 7-azabicyclo[4.4.1]undecyl and 8-azabicyclo[5.4.1]dodecyl, especially 5-azabicyclo[2.2.1]heptyl and 6-azabicyclo[3.2.1]octyl.


Where  $R^7$  represents a  $C_{2-4}$ alkyl group substituted by a 5 or 6 membered heteroaliphatic ring containing one or two heteroatoms selected from N, O and S, suitable rings include pyrrolidino, piperidino, piperazino,

morpholino, or thiomorpholino. Particularly preferred are nitrogen containing heteroaliphatic rings, especially pyrrolidino and morpholino rings.


Particularly apt values for X for compounds of the formulae (I) or (Ia) include  $\text{CH}_2$ ,  $\text{CH}(\text{CH}_3)$  and  $\text{CH}_2\text{CH}_2$  of which the  $\text{CH}_2$  group is preferred.

Favourably Het is a 5-membered ring.


In particular, Het may represent a heterocyclic ring selected from:




Particularly preferred heterocyclic rings represented by  $\text{Het-ZN}(\text{O})_m\text{R}^6\text{R}^7$  are selected from:



Most especially,  $\text{Het-ZN}(\text{O})_m\text{R}^6\text{R}^7$  may represent a heterocyclic ring selected from:




A particularly preferred heterocyclic ring represented by  
Het-ZN(O)<sub>m</sub>R<sup>6</sup>R<sup>7</sup> is:



5

One favoured group of compounds of this invention are of the  
formula (Ib) and pharmaceutically acceptable salts thereof:




10

wherein A<sup>1</sup>, A<sup>2</sup> and A<sup>3</sup> are defined in relation to formula (Ia) and wherein  
Z, R<sup>6</sup>, R<sup>7</sup>, m and n are as defined in relation to formula (I).

Another favoured group of compounds of the present invention are  
of the formula (Ic) and pharmaceutically acceptable salts thereof:

15



wherein  $A^1$ ,  $A^2$  and  $A^3$  are defined in relation to formula (Ia),  $Q^2$  is CH or N and Z,  $R^6$ ,  $R^7$ , m and n are as defined in relation to formula (I).

5 With respect to compounds of the formulae (I), (Ia), (Ib) and (Ic), Z may be a linear, branched or cyclic group. Favourably Z contains 1 to 4 carbon atoms and most favourably 1 or 2 carbon atoms. A particularly favourable group Z is  $CH_2$ .

10 With respect to compounds of the formulae (I), (Ia), (Ib) and (Ic),  $R^6$  may aptly be a  $C_{1-4}$ alkyl group or a  $C_{2-4}$ alkyl group substituted by a hydroxyl or  $C_{1-2}$ alkoxy group,  $R^7$  may aptly be a  $C_{1-4}$ alkyl group or a  $C_{1-4}$ alkyl group substituted by a hydroxyl or  $C_{1-2}$ alkoxy group, or  $R^6$  and  $R^7$  may be linked so that, together with the nitrogen atom to which they are attached, they form an azetidiny, pyrrolidiny, piperidyl, morpholino, 15 thiomorpholino, piperazino or piperazino group substituted on the nitrogen atom by a  $C_{1-4}$ alkyl group or a  $C_{2-4}$ alkyl group substituted by a hydroxy or  $C_{1-2}$ alkoxy group.


Particularly suitable moieties  $ZNR^6R^7$  include those wherein Z is  $CH_2$  or  $CH_2CH_2$  and  $NR^6R^7$  is amino, methylamino, dimethylamino, 20 diethylamino, azetidiny, pyrrolidino and morpholino.

Further preferred moieties represented by  $ZNR^6R^7$  are those wherein Z is  $CH_2$  or  $CH_2CH_2$ ,  $R^6$  represents hydrogen,  $C_{1-4}$ alkyl or  $C_{3-6}$ cycloalkyl and  $R^7$  is  $C_{2-4}$ alkyl substituted by one or two substituents

selected from hydroxy, C<sub>1-2</sub>alkoxy, azetidiny, pyrrolidino, piperidino, morpholino or thiomorpholino.

In particular, Z is preferably CH<sub>2</sub> and NR<sup>6</sup>R<sup>7</sup> is preferably dimethylamino, azetidiny or pyrrolidino, especially dimethylamino.

5 According to a further aspect of the present invention, or particularly preferred class of compound is that represented by formula (Id) and pharmaceutically acceptable salts thereof:



10

wherein m and n are as defined in relation to formula (I).

Specific compounds within the scope of the present invention include:

- 15 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(N,N-dimethylamino(N-oxide)methyl)-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine;
- 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(N,N-dimethylaminomethyl)-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine N-oxide;
- 20 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(N,N-dimethylamino(N-oxide)methyl)-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine N-oxide;



3-(S)-(4-fluorophenyl)-2-(R)-(1-(R)-(3-methylthio-5-(trifluoromethyl) phenyl)ethoxy)-4-(5-pyrrolidinomethyl-1,2,3-triazol-4-yl)methylmorpholine N-oxide;

3-(S)-(4-fluorophenyl)-2-(R)-(1-(R)-(3-methylthio-5-(trifluoromethyl) phenyl)ethoxy)-4-(5-pyrrolidino(N-oxide)methyl-1,2,3-triazol-4-yl)methylmorpholine N-oxide;

and pharmaceutically acceptable salts thereof.

For use in medicine, the salts of the compounds of formula (I) will be non-toxic pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds according to the invention or of their non-toxic pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts such as those formed with hydrochloric acid, fumaric acid, p-toluenesulphonic acid, maleic acid, succinic acid, acetic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Salts of amine groups may also comprise quaternary ammonium salts in which the amino nitrogen atom carries a suitable organic group such as an alkyl, alkenyl, alkynyl or aralkyl moiety. Furthermore, where the compounds of the invention carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include metal salts such as alkali metal salts, e.g. sodium or potassium salts; and alkaline earth metal salts, e.g. calcium or magnesium salts.

The pharmaceutically acceptable salts of the present invention may be formed by conventional means, such as by reacting the free base form of the product with one or more equivalents of the appropriate acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is removed *in vacuo* or by freeze drying or by exchanging the anions of an existing salt for another anion on a suitable ion exchange resin.

The present invention includes within its scope prodrugs of the compounds of formula (I) above. In general, such prodrugs will be

functional derivatives of the compounds of formula (I) which are readily convertible *in vivo* into the required compound of formula (I).


Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs",  
5 ed. H. Bundgaard, Elsevier, 1985.

A prodrug may be a pharmacologically inactive derivative of a biologically active substance (the "parent drug" or "parent molecule") that requires transformation within the body in order to release the active  
10 drug, and that has improved delivery properties over the parent drug molecule. The transformation *in vivo* may be, for example, as the result of some metabolic process, such as chemical or enzymatic hydrolysis of a carboxylic, phosphoric or sulphate ester, or reduction or oxidation of a susceptible functionality.

The present invention includes within its scope solvates of the  
15 compounds of formula (I) and salts thereof, for example, hydrates.

The compounds according to the invention have at least three asymmetric centres, and may accordingly exist both as enantiomers and as diastereoisomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present  
20 invention.

The preferred compounds of the formula (I), (Ia), (Ib), (Ic) and (Id) will have the 2- and 3- substituent cis and the preferred stereochemistry at the 2-position is that possessed by the compound of Example 1 (i.e. 2-(R)-), the preferred stereochemistry of the 3-position is that possessed by  
25 the compound of Example 1 (i.e. 3-(S)), and the preferred stereochemistry of the carbon to which the group Y is attached is either (R) when Y is C<sub>1-4</sub>alkyl (e.g. methyl) or (S) when Y is C<sub>1-4</sub>alkyl substituted by hydroxy (e.g. CH<sub>2</sub>OH). Thus for example as shown in formula (Ie)



The present invention further provides pharmaceutical compositions comprising one or more compounds of formula (I) in association with a pharmaceutically acceptable carrier.

Preferably the compositions according to the invention are in unit dosage forms such as tablets, pills, capsules, powders, granules, solutions or suspensions, or suppositories, for oral, parenteral or rectal administration, or administration by inhalation or insufflation.

For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose, sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a non-toxic pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The tablets or pills of the novel composition can be coated or otherwise compounded to provide a

dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to  
5 resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

10 The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar  
15 pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

Preferred compositions for administration by injection include those  
20 comprising a compound of formula (I), as the active ingredient, in association with a surface-active agent (or wetting agent or surfactant) or in the form of an emulsion (as a water-in-oil or oil-in-water emulsion).

Suitable surface-active agents include, in particular, non-ionic agents, such as polyoxyethylenesorbitans (e.g. Tween™ 20, 40, 60, 80 or  
25 85) and other sorbitans (e.g. Span™ 20, 40, 60, 80 or 85). Compositions with a surface-active agent will conveniently comprise between 0.05 and 5% surface-active agent, and preferably between 0.1 and 2.5%. It will be appreciated that other ingredients may be added, for example mannitol or other pharmaceutically acceptable vehicles, if necessary.

30 Suitable emulsions may be prepared using commercially available fat emulsions, such as Intralipid™, Liposyn™, Infonutrol™, Lipofundin™

and Lipiphysan™. The active ingredient may be either dissolved in a pre-mixed emulsion composition or alternatively it may be dissolved in an oil (e.g. soybean oil, safflower oil, cottonseed oil, sesame oil, corn oil or almond oil) and an emulsion formed upon mixing with a phospholipid (e.g. egg phospholipids, soybean phospholipids or soybean lecithin) and water. It will be appreciated that other ingredients may be added, for example glycerol or glucose, to adjust the tonicity of the emulsion. Suitable emulsions will typically contain up to 20% oil, for example, between 5 and 20%. The fat emulsion will preferably comprise fat droplets between 0.1 and 1.0µm, particularly 0.1 and 0.5µm, and have a pH in the range of 5.5 to 8.0.

Particularly preferred emulsion compositions are those prepared by mixing a compound of formula (I) with Intralipid™ or the components thereof (soybean oil, egg phospholipids, glycerol and water).

Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as set out above. Preferably the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions in preferably sterile pharmaceutically acceptable solvents may be nebulised by use of inert gases. Nebulised solutions may be breathed directly from the nebulising device or the nebulising device may be attached to a face mask, tent or intermittent positive pressure breathing machine. Solution, suspension or powder compositions may be administered, preferably orally or nasally, from devices which deliver the formulation in an appropriate manner.

The present invention further provides a process for the preparation of a pharmaceutical composition comprising a compound of formula (I), which process comprises bringing a compound of formula (I) into association with a pharmaceutically acceptable carrier or excipient.

The compounds of formula (I) are of value in the treatment of a wide variety of clinical conditions which are characterised by the presence of an excess of tachykinin, in particular substance P, activity. These may include disorders of the central nervous system such as anxiety,

5 depression, psychosis and schizophrenia; epilepsy; neurodegenerative disorders such as dementia, including AIDS related dementia, senile dementia of the Alzheimer type, Alzheimer's disease and Down's syndrome; demyelinating diseases such as multiple sclerosis (MS) and amyotrophic lateral sclerosis (ALS; Lou Gehrig's disease) and other

10 neuropathological disorders such as peripheral neuropathy, for example AIDS related neuropathy, diabetic and chemotherapy-induced neuropathy, and postherpetic and other neuralgias; neuronal damage, such as cerebral ischemic damage and cerebral edema in cerebrovascular disorders; small cell carcinomas such as small cell lung cancer; respiratory

15 diseases, particularly those associated with excess mucus secretion such as chronic obstructive airways disease, bronchopneumonia, chronic bronchitis, asthma, and bronchospasm; airways diseases modulated by neurogenic inflammation; diseases characterised by neurogenic mucus secretion, such as cystic fibrosis; diseases associated with decreased

20 glandular secretions, including lacrimation, such as Sjogren's syndrome, hyperlipoproteinemias IV and V, hemochromatosis, sarcoidosis, and amyloidosis; inflammatory diseases such as inflammatory bowel disease, psoriasis, fibrositis, ocular inflammation, osteoarthritis, rheumatoid arthritis, pruritis and sunburn; allergies such as eczema and rhinitis;

25 hypersensitivity disorders such as poison ivy; ophthalmic diseases such as conjunctivitis, vernal conjunctivitis, dry eye syndrome, and the like; ophthalmic conditions associated with cell proliferation such as proliferative vitreoretinopathy; cutaneous diseases such as contact dermatitis, atopic dermatitis, urticaria, and other eczematoid dermatitis;

30 addiction disorders including the withdrawal response produced by chronic treatment with, or abuse of, drugs such as benzodiazepines,

opiates, cocaine, alcohol and nicotine; stress related somatic disorders; reflex sympathetic dystrophy such as shoulder/hand syndrome; dysthymic disorders; adverse immunological reactions such as rejection of transplanted tissues and disorders related to immune enhancement or suppression such as systemic lupus erythematosus; gastrointestinal (GI) disorders, including inflammatory disorders, and diseases of the GI tract, such as gastritis, gastroduodenal ulcers, gastric carcinomas, gastric lymphomas, disorders associated with the neuronal control of viscera, ulcerative colitis, Crohn's disease, irritable bowel syndrome and emesis, including acute, delayed, post-operative, late phase or anticipatory emesis such as emesis induced by chemotherapy, radiation, toxins, viral or bacterial infections, pregnancy, vestibular disorders, motion, surgery, migraine, opioid analgesics, and variations in intracranial pressure, in particular, for example, drug or radiation induced emesis or post-operative nausea and vomiting; disorders of bladder function such as cystitis, bladder detrusor hyper-reflexia and incontinence; fibrosing and collagen diseases such as scleroderma and eosinophilic fasciitis; disorders of blood flow caused by vasodilation and vasospastic diseases such as angina, migraine and Reynaud's disease; and pain or nociception, for example, that attributable to or associated with any of the foregoing conditions, especially the transmission of pain in migraine.

Hence, the compounds of the present invention may be of use in the treatment of physiological disorders associated with excessive stimulation of tachykinin receptors, especially neurokinin-1 receptors, and as neurokinin-1 antagonists for the control and/or treatment of any of the aforementioned clinical conditions in mammals, including humans.

The compounds of formula (I) are also of value in the treatment of a combination of the above conditions, in particular in the treatment of combined post-operative pain and post-operative nausea and vomiting.

The compounds of formula (I) are particularly useful in the treatment of emesis, including acute, delayed, post-operative, late phase or

anticipatory emesis, such as emesis or nausea induced by chemotherapy, radiation, toxins, such as metabolic or microbial toxins, viral or bacterial infections, pregnancy, vestibular disorders, motion, mechanical stimulation, gastrointestinal obstruction, reduced gastrointestinal motility, visceral pain, psychological stress or disturbance, high altitude, weightlessness, opioid analgesics, intoxication, resulting for example from consumption of alcohol, surgery, migraine, and variations in intracranial pressure. Most especially, the compounds of formula (I) are of use in the treatment of emesis induced by antineoplastic (cytotoxic) agents including those routinely used in cancer chemotherapy.

Examples of such chemotherapeutic agents include alkylating agents, for example, nitrogen mustards, ethyleneimine compounds, alkyl sulphonates and other compounds with an alkylating action such as nitrosoureas, cisplatin and dacarbazine; antimetabolites, for example, folic acid, purine or pyrimidine antagonists; mitotic inhibitors, for example, vinca alkaloids and derivatives of podophyllotoxin; and cytotoxic antibiotics.

Particular examples of chemotherapeutic agents are described, for instance, by D. J. Stewart in *"Nausea and Vomiting: Recent Research and Clinical Advances"*, Eds. J. Kuucharczyk *et al*, CRC Press Inc., Boca Raton, Florida, USA (1991) pages 177-203, especially page 188.

Commonly used chemotherapeutic agents include cisplatin, dacarbazine (DTIC), dactinomycin, mechlorethamine (nitrogen mustard), streptozocin, cyclophosphamide, carmustine (BCNU), lomustine (CCNU), doxorubicin (adriamycin), daunorubicin, procarbazine, mitomycin, cytarabine, etoposide, methotrexate, 5-fluorouracil, vinblastine, vincristine, bleomycin and chlorambucil [R. J. Gralla *et al* in *Cancer Treatment Reports* (1984) 68(1), 163-172].

The compounds of formula (I) are also of use in the treatment of emesis induced by radiation including radiation therapy such as in the



treatment of cancer, or radiation sickness; and in the treatment of post-operative nausea and vomiting.

It will be appreciated that the compounds of formula (I) may be presented together with another therapeutic agent as a combined  
5 preparation for simultaneous, separate or sequential use for the relief of emesis. Such combined preparations may be, for example, in the form of a twin pack.

A further aspect of the present invention comprises the compounds of formula (I) in combination with a 5-HT<sub>3</sub> antagonist, such as  
10 ondansetron, granisetron or tropisetron, or other anti-emetic medicaments, for example, a dopamine antagonist such as metoclopramide or GABA<sub>A</sub> receptor agonists such as baclofen. Additionally, a compound of formula (I) may be administered in combination with an anti-  
inflammatory corticosteroid, such as dexamethasone, triamcinolone,  
15 triamcinolone acetonide, flunisolide, budesonide, or others such as those disclosed in US patent nos. 2,789,118, 2,990,401, 3,048,581, 3,126,375, 3,929,768, 3,996,359, 3,928,326 and 3,749,712. Dexamethasone (Decadron™) is particularly preferred. Furthermore, a compound of formula (I) may be administered in combination with a chemotherapeutic  
20 agent such as an alkylating agent, antimetabolite, mitotic inhibitor or cytotoxic antibiotic, as described above. In general, the currently available dosage forms of the known therapeutic agents for use in such combinations will be suitable.

When tested in the ferret model of cisplatin-induced emesis  
25 described by F. D. Tattersall *et al*, in *Eur. J. pharmacol.*, (1993) 250, R5-R6, the compounds of the present invention were found to attenuate the retching and vomiting induced by cisplatin.

The compounds of formula (I) are also particularly useful in the treatment of pain or nociception and/or inflammation and disorders  
30 associated therewith such as, for example, neuropathy, such as diabetic and chemotherapy-induced neuropathy, postherpetic and other

neuralgias, asthma, osteoarthritis, rheumatoid arthritis, headache and especially migraine.

The present invention further provides a compound of formula (I) for use in therapy.

5       According to a further or alternative aspect, the present invention provides a compound of formula (I) for use in the manufacture of a ~~medicament for the treatment of physiological disorders associated with~~

---

an excess of tachykinins, especially substance P.

10       The present invention also provides a method for the the treatment or prevention of physiological disorders associated with an excess of tachykinins, especially substance P, which method comprises administration to a patient in need thereof of a tachykinin reducing amount of a compound of formula (I) or a composition comprising a compound of formula (I).

15       For the treatment of certain conditions it may be desirable to employ a compound according to the present invention in conjunction with another pharmacologically active agent. For example, for the treatment of respiratory diseases such as asthma, a compound of formula (I) may be used in conjunction with a bronchodilator, such as a  $\beta_2$ -adrenergic receptor  
20 agonist or tachykinin antagonist which acts at NK-2 receptors. The compound of formula (I) and the bronchodilator may be administered to a patient simultaneously, sequentially or in combination.

Likewise, a compound of the present invention may be employed with a leukotriene antagonists, such as a leukotriene D<sub>4</sub> antagonist such  
25 as a compound selected from those disclosed in European patent specification nos. 0 480 717 and 0 604 114 and in US patent nos. 4,859,692 and 5,270,324. This combination is particularly useful in the treatment of respiratory diseases such as asthma, chronic bronchitis and cough.

The present invention accordingly provides a method for the  
30 treatment of a respiratory disease, such as asthma, which method comprises administration to a patient in need thereof of an effective

amount of a compound of formula (I) and an effective amount of a bronchodilator.

The present invention also provides a composition comprising a compound of formula (I), a bronchodilator, and a pharmaceutically acceptable carrier.

It will be appreciated that for the treatment or prevention of migraine, a compound of the present invention may be used in conjunction with other anti-migraine agents, such as ergotamines or 5-HT<sub>1</sub> agonists, especially sumatriptan.

Likewise, for the treatment of behavioural hyperalgesia, a compound of the present invention may be used in conjunction with an antagonist of N-methyl D-aspartate (NMDA), such as dizocilpine.

For the treatment or prevention of inflammatory conditions in the lower urinary tract, especially cystitis, a compound of the present invention may be used in conjunction with an antiinflammatory agent such as a bradykinin receptor antagonist.


In the treatment of the conditions associated with an excess of tachykinins, a suitable dosage level is about 0.001 to 50 mg/kg per day, in particular about 0.01 to about 25 mg/kg, such as from about 0.05 to about 10 mg/kg per day.

For example, in the treatment of conditions involving the neurotransmission of pain sensations, a suitable dosage level is about 0.001 to 25 mg/kg per day, preferably about 0.005 to 10 mg/kg per day, and especially about 0.005 to 5 mg/kg per day. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day.

In the treatment of emesis using an injectable formulation, a suitable dosage level is about 0.001 to 10 mg/kg per day, preferably about 0.005 to 5 mg/kg per day, and especially 0.01 to 1 mg/kg per day. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day.

It will be appreciated that the amount of a compound of formula (I) required for use in any treatment will vary not only with the particular compounds or composition selected but also with the route of administration, the nature of the condition being treated, and the age and condition of the patient, and will ultimately be at the discretion of the attendant physician.

According to one general process, the compounds of formula (I) may be prepared from compounds of formula (II)




by oxidation of one or both of the nitrogen atoms drawn in formula (II).

The oxidation reaction may be effected using hydrogen peroxide, an alkyl peroxide such as t-butylperoxide or a peroxy acid (peracid) such as m-chloroperbenzoic acid or peracetic acid. Hydrogen peroxide is particularly preferred. The reaction is conveniently effected at room temperature in a solvent such as an ether, for example, tetrahydrofuran, acetonitrile or a chlorinated hydrocarbon, for example, dichloromethane, or an oil, for example, soya bean oil.

Where oxidation of the morpholine nitrogen is desired, the reaction is conveniently effected in acidic conditions, for example, using hydrogen peroxide in acetic acid.

Where oxidation of the amine moiety  $\text{NR}^6\text{R}^7$  is desired, neutral conditions are preferred, for example, using hydrogen peroxide in soya bean oil.

Compounds of formula (II) may be prepared by a variety of methods, thus, according to intermediate process (A), the compounds of formula (II) may be prepared from compounds of formula (IIIa)




wherein  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^4$ ,  $\text{R}^5$  and  $\text{Y}$  are as defined in relation to formula (I) by reaction with a compound of formula (IIIb):



where  $\text{X}$  is as defined in relation to formula (I),  $\text{Het}'$  is as defined for the group  $\text{Het}$  in relation to formula (I) or a precursor therefor and  $\text{L}$  is a leaving group for example a halogen atom such as bromine or chlorine; and, if  $\text{Het}'$  is a precursor group, converting it to a group  $\text{Het}$  (in which process any reactive group may be protected and thereafter deprotected if desired).


This reaction may be performed in conventional manner, for example in an organic solvent such as dimethylformamide in the presence of an acid acceptor such as potassium carbonate.

According to another process (B), compounds of formula (II) wherein  $\text{Het}$  represents 1,2,3-triazol-4-yl substituted by  $\text{CH}_2\text{NR}^6\text{R}^7$ , and  $\text{X}$  is  $-\text{CH}_2-$ , may be prepared by reaction of a compound of formula (IV)



- with an azide, for example, sodium azide in a suitable solvent such as
- 5 dimethylsulphoxide at a temperature of between 40°C and 100°C, followed by reduction of the carbonyl group adjacent to -NR<sup>6</sup>R<sup>7</sup> using a suitable reducing agent such as lithium aluminium hydride at a temperature between -10°C and room temperature, conveniently at room temperature.


- Alternatively, according to a process (C), compounds of formula (II)
- 10 wherein Het represents 1,2,3-triazol-4-yl substituted by CH<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, and X is -CH<sub>2</sub>-, may be prepared by reaction of a compound of formula (V)



- 15 with an amine of formula NHR<sup>6</sup>R<sup>7</sup>, in a suitable solvent such as an ether, for example, dioxan, at elevated temperature, for example, between 50°C

and 100°C, in a sealed tube, or the like. This reaction is based upon that described in *Chemische Berichte* (1989) 122, p. 1963.



According to another process, (D), compounds of formula (II) wherein Het represents 1,3,5-triazine may be prepared by reaction of  
5 intermediates of formula (VI):



with an appropriately substituted 1,3,5-triazine.

The reaction is conveniently effected in a suitable organic solvent, such as acetonitrile, at elevated temperature, such as 80-90°C, preferably  
10 about 82°C.

According to a further process, (E), compounds of formula (II) wherein Het represents 1,2,4-triazine may be prepared by reaction of an intermediate of formula (VII) with a dicarbonyl compound of formula  
(VIII):




15

wherein R<sup>35</sup> represents the substituent ZNR<sup>6</sup>R<sup>7</sup>.

The reaction is conveniently effected in a suitable organic solvent, such as an ether, e.g. tetrahydrofuran, conveniently at ambient temperature.

- 5 According to a further process (F), compounds of formula (II) wherein Het represents a substituted 1,2,4-triazolyl group may be prepared by reaction of an intermediate of formula (III) with a compound of formula (IX)



(IX)


- 10 wherein X is as defined in relation to formula (I), Hal is a halogen atom, for example, bromine, chlorine or iodine and R<sup>18</sup> is H, CONH<sub>2</sub> or OCH<sub>3</sub> (which is converted to an oxo substituent under the reaction conditions), in the presence of a base, followed where necessary by conversion to a compound of formula (I), for example, by reduction of the CONH<sub>2</sub> group to CH<sub>2</sub>NH<sub>2</sub>.

- 15 Suitable bases of use in the reaction include alkali metal carbonates such as, for example, potassium carbonate. The reaction is conveniently effected in an anhydrous organic solvent such as, for example, anhydrous dimethylformamide, preferably at elevated temperature, such as about 140°C.

- 20 A suitable reducing agent for the group CONH<sub>2</sub> is lithium aluminium hydride, used at between -10°C and room temperature.

According to another process, (G), compounds of formula (II) wherein Het represents thioxotriazolyl may be prepared from intermediates of formula (X)





(X)


by reaction with a compound of formula  $\text{HNCS}$ , in the presence of a base.

Suitable bases of use in the reaction include organic bases such as, for example, 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU). The reaction is conveniently effected in a suitable organic solvent, such as alcohol, e.g. butanol.

Further details of suitable procedures will be found in the accompanying Examples.

Intermediates of formula (IV) may be prepared from intermediates of formula (IIIa) by reaction with an acetylene compound of formula  $\text{HC}\equiv\text{C}-\text{CH}_2-\text{Hal}$  in the presence of a base such as potassium carbonate in a suitable solvent such as dimethylformamide, conveniently at room temperature, followed by reaction of the resultant acetylene intermediate with an amide of formula  $\text{Hal}-\text{CO}-\text{NR}^6\text{R}^7$  in the presence of suitable catalysts including bis(triphenylphosphine) palladium(II) chloride, copper(I) iodide and triphenylphosphine in a suitable solvent such as triethylamine, preferably at reflux.

Intermediates of formula (V) may be prepared from a compound of formula (XI)



(XI)

wherein Hal is a halogen atom, for example, chlorine, bromine or iodine, especially chlorine, by reaction with an azide, for example, sodium azide in a suitable solvent such as dimethylsulphoxide at or below room temperature.

Compounds of formula (XI) may be prepared by a dropwise addition of an intermediate of formula (IIIa) to a dihaloacetylene of formula Hal-CH<sub>2</sub>-C≡C-CH<sub>2</sub>-Hal where each Hal is independently chlorine, bromine or iodine, especially chlorine. The reaction is conveniently effected in a suitable solvent such as dimethylformamide in the presence of a base such as potassium carbonate.

Intermediates of formula (VI) may be prepared from intermediates of formula (IIIa) by reaction with a compound of formula Hal-X-C(NH)NH<sub>2</sub>, where Hal and X are as previously defined.

Intermediates of formula (VII) may be prepared from intermediates of formula (IIIa) by reaction with a compound of formula Hal-X-C(NH)NHNH-Boc, wherein Hal and X are as previously defined and Boc stands for t-butoxycarbonyl, followed by deprotection under acidic conditions.

Compounds of formula (VIII) are commercially available or may be prepared from commercially available compounds by known methods.

Compounds of formula (IX) may be prepared as described in *J. Med. Chem.*, (1984) 27, 849.


Intermediates of formula (X) may be prepared from the corresponding ester by treatment with hydrazine. The reaction is conveniently effected in a suitable organic solvent, such as an alcohol, for example, ethanol, at elevated temperature.

5 For compounds wherein Het is a heterocycle bearing the  $\text{ZNR}^6\text{R}^7$  group where Z is  $\text{CH}_2$ , certain favoured compounds of formula (I) may be prepared from a corresponding compound with a hydrogen atom in place of the  $\text{ZNR}^6\text{R}^7$ . Thus, for example a compound of the formula (I) wherein  
10 Het is an imidazolinone group carrying a  $\text{CH}_2\text{NR}^6\text{R}^7$  moiety may be prepared from a corresponding compound lacking the  $\text{CH}_2\text{NR}^6\text{R}^7$  moiety by reaction with formaldehyde and an amine  $\text{NHR}^6\text{R}^7$  under conventional Mannich reaction conditions, for example in methanol with heating. If  
15 desired a pre-formed reagent such as  $\text{R}^6\text{R}^7\text{N}^+=\text{CH}_2.\text{I}^-$  may be employed and a tertiary amine such as triethylamine used as acid acceptor.

15 Alternatively a compound of formula (I) wherein Het is an imidazolinone group lacking a  $\text{CH}_2\text{NR}^6\text{R}^7$  may be reacted with paraformaldehyde and an amine for example a secondary amine such as pyrrolidine to give a compound wherein the imidazolinone ring is substituted by  $\text{CH}_2\text{NR}^6\text{R}^7$  where  $\text{R}^6$ ,  $\text{R}^7$  and the nitrogen atom to which  
20 they are attached form a heteroaliphatic ring of 4 to 7 ring atoms which may optionally contain an oxygen ring atom or a second nitrogen atom which will be part of a  $\text{NR}^8$  moiety, where  $\text{R}^8$  is as previously defined.

This reaction may be performed in a conventional manner, for instance, in a suitable solvent such as an alcohol, for example, methanol  
25 at an elevated temperature up to the boiling point of the solvent.

A further alternative method for the preparation of certain compounds of formula (II) involves the reaction of an intermediate of formula (IIIa) as defined above with one of the compounds of formula (XII):




wherein each LG, which may be the same or different, is a leaving group, such as an alkyl- or arylsulphonyloxy group (e.g. mesylate or tosylate) or, in particular, a halogen atom, (e.g. bromine, chlorine or iodine) and X and Z are as defined in formula (I), followed by reaction of the resultant compound with an amine  $\text{NHR}^6\text{R}^7$  to complete the  $\text{ZNR}^6\text{R}^7$  moiety.

This reaction is conveniently effected in an organic solvent such as dimethylformamide in the presence of an acid acceptor such as potassium carbonate.

It will be appreciated that, where necessary, reactive groups may be protected, thus for example, the NH groups of an imidazolinone of formula (XIIa) may be protected by any suitable amine protecting group such as an acetyl group.

The compounds of the formula (IIIa) may be prepared as shown in the following Scheme in which  $\text{Ar}^1$  represents the  $\text{R}^1, \text{R}^2, \text{R}^3$  substituted phenyl group;  $\text{Ar}^2$  represents the  $\text{R}^4, \text{R}^5$  substituted phenyl group and Ph represents phenyl:



L-Selectride is lithium tri-sec-butylborohydride.

The following references describe methods which may be applied by the skilled worker to the chemical synthesis set forth above once the skilled worker has read the disclosure herein:

- (i) D.A. Evans *et al.*, *J. Am. Chem. Soc.*, (1990) 112, 4011.
- (ii) I. Yanagisawa *et al.*, *J. Med. Chem.*, (1984) 27, 849.
- (iii) R. Duschinsky *et al.*, *J. Am. Chem. Soc.*, (1948) 70, 657.
- (iv) F.N. Tebbe *et al.*, *J. Am. Chem. Soc.*, (1978) 100, 3611.
- (v) N.A. Petasis *et al.*, *J. Am. Chem. Soc.*, (1990) 112, 6532.
- (vi) K. Takai *et al.*, *J. Org. Chem.*, (1987) 52, 4412.

The Examples disclosed herein produce predominantly the preferred isomers. The unfavoured isomers are also produced as minor components. If desired they may be isolated and employed to prepare the

various stereoisomers in conventional manner, for example chromatography using an appropriate column. However, the skilled worker will appreciate that although the Examples have been optimized to the production of the preferred isomers, variation in solvent, reagents, chromatography etc can be readily employed to yield the other isomers.

It will be appreciated that compounds of the formula (I) wherein Het contains an =O or =S substituent can exist in tautomeric forms. All such tautomeric forms and mixtures thereof are included within this invention. Most aptly the =O or =S substituent in Het is the =O substituent.

Where they are not commercially available, the intermediates of formula (III) above may be prepared by the procedures described in the accompanying Examples or by alternative procedures which will be readily apparent to one skilled in the art.

During any of the above synthetic sequences it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *Protective Groups in Organic Chemistry*, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

The exemplified compounds of this invention were tested by the methods set out at pages 36 to 39 of International Patent Specification No. WO 93/01165. The compounds or, in the case of prodrugs, the parent compounds, were found to be active with  $IC_{50}$  at the  $NK_1$  receptor of less than 100nM on said test method.

The following Examples illustrate the preparation of compounds according to the present invention:

DESCRIPTION 1(S)-(4-Fluorophenyl)glycineVia Chiral Synthesis:5    Step A: 3-(4-Fluorophenyl)acetyl-4-(S)-benzyl-2-oxazolidinone

An oven-dried, 1 L 3-necked flask, equipped with a septum, nitrogen inlet, thermometer, and a magnetic stirring bar, was flushed with nitrogen and charged with a solution of 5.09g (33.0mmol) of 4-fluorophenylacetic acid in 100ml of anhydrous ether. The solution was cooled to -10°C and treated with 5.60ml (40.0mmol) of triethylamine followed by 4.30ml (35.0mmol) of trimethylacetyl chloride. A white precipitate formed immediately. The resulting mixture was stirred at -10°C for 40 minutes, then cooled to -78°C.

15    An oven-dried, 250ml round bottom flask, equipped with a septum and a magnetic stirring bar, was flushed with nitrogen and charged with a solution of 5.31g (30.0mmol) of 4-(S)-benzyl-2-oxazolidinone in 40ml of dry THF. The solution was stirred in a dry ice/acetone bath for 10 minutes, then 18.8ml of 1.6M n-butyllithium solution in hexanes was slowly added. After 10 minutes, the lithiated oxazolidinone solution was added, via  
20    cannula, to the above mixture in the 3-necked flask. The cooling bath was removed from the resulting mixture and the temperature was allowed to rise to 0°C. The reaction was quenched with 100ml of saturated aqueous ammonium chloride solution, transferred to a 1l flask, and the ether and THF were removed *in vacuo*. The concentrated mixture was partitioned  
25    between 300ml of methylene chloride and 50ml of water and the layers were separated. The organic layer was washed with 100ml of 2N aqueous hydrochloric acid solution, 300ml of saturated aqueous sodium bicarbonate solution, dried over magnesium sulfate and concentrated *in vacuo*. Flash chromatography on 400g of silica gel using 3:2 v/v  
30    hexanes/ether as the eluant afforded 8.95g of an oil that slowly solidified on standing. Recrystallisation from 10:1 hexanes/ether afforded 7.89g

(83%) of the title compound as a white solid: mp 64-66°C. MS (FAB):  $m/z$  314 ( $M^+ + H$ , 100%), 177 ( $M - ArCH_2CO + H$ , 85%).  $^1H$  NMR (400MHz,  $CDCl_3$ )  $\delta$  2.76 (1H, dd,  $J=13.2, 9.2$ Hz), 3.26 (dd,  $J=13.2, 3.2$ Hz), 4.16-4.34 (4H, m), 4.65 (1H, m), 7.02-7.33 (9H, m).

- 5 Analysis Calcd. for  $C_{18}H_{18}FNO_3$ : C, 69.00; H, 5.15; N, 4.47; F, 6.06;  
Found: C, 68.86; H, 5.14; N, 4.48; F, 6.08%.

Step B: 3-((S)-Azido-(4-fluorophenyl))acetyl-4-(S)-benzyl-2-oxazolidinone

- An oven-dried, 1l 3-necked flask, equipped with a septum, nitrogen  
10 inlet, thermometer, and a magnetic stirring bar, was flushed with nitrogen  
and charged with a solution of 58.0ml of 1M potassium  
bis(trimethylsilyl)amide solution in toluene and 85ml of THF and was  
cooled to -78°C. An oven-dried 250ml round-bottomed flask, equipped  
with a septum and a magnetic stirring bar, was flushed with nitrogen and  
15 charged with a solution of 7.20g (23.0mmol) of 3-(4-fluorophenyl)acetyl-4-  
(S)-benzyl-2-oxazolidinone (from Step A) in 40ml of THF. The acyl  
oxazolidinone solution was stirred in a dry ice/acetone bath for 10  
minutes, then transferred, via cannula, to the potassium  
bis(trimethylsilyl)amide solution at such a rate that the internal  
20 temperature of the mixture was maintained below -70°C. The acyl  
oxazolidinone flask was rinsed with 15ml of THF and the rinse was added,  
via cannula, to the reaction mixture and the resulting mixture was stirred  
at -78°C for 30 minutes. An oven-dried, 250ml round-bottomed flask,  
equipped with a septum and a magnetic stirring bar, was flushed with  
25 nitrogen and charged with a solution of 10.89g (35.0mmol) of  
2,4,6-triisopropylphenylsulfonyl azide in 40ml of THF. The azide solution  
was stirred in a dry ice/acetone bath for 10 minutes, then transferred, via  
cannula, to the reaction mixture at such a rate that the internal  
temperature of the mixture was maintained below -70°C. After 2 minutes,  
30 the reaction was quenched with 6.0ml of glacial acetic acid, the cooling  
bath was removed and the mixture was stirred at room temperature for 18



hours. The quenched reaction mixture was partitioned between 300ml of ethyl acetate and 300ml of 50% saturated aqueous sodium bicarbonate solution. The organic layer was separated, dried over magnesium sulfate, and concentrated *in vacuo*. Flash chromatography on 500g of silica gel using 2:1 v/v, then 1:1 v/v hexanes/methylene chloride as the eluant afforded 5.45g (67%) of the title compound as an oil. IR Spectrum (neat,  $\text{cm}^{-1}$ ): 2104, 1781, 1702.  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ )  $\delta$  2.86 (1H, dd,  $J=13.2$ , 9.6Hz), 3.40 (1H, dd,  $J=13.2$ , 3.2Hz), 4.09-4.19 (2H, m), 4.62-4.68 (1H, m), 6.14 (1H, s), 7.07-7.47 (9H, m).

Analysis Calcd. for  $\text{C}_{18}\text{H}_{15}\text{FN}_4\text{O}_3$ : C 61.01; H, 4.27; N, 15.81; F, 5.36; Found: C, 60.99; H, 4.19; N, 15.80; F, 5.34%.

Step C: (S)-Azido-(4-fluorophenyl)acetic acid

A solution of 5.40g (15.2mmol) of 3-((S)-azido-(4-fluorophenyl))acetyl-4-(S)-benzyl-2-oxazolidinone (from Step B) in 200ml of 3:1 v/v THF/water was stirred in an ice bath for 10 minutes. 1.28g (30.4mmol) of lithium hydroxide monohydrate was added in one portion and the resulting mixture was stirred cold for 30 minutes. The reaction mixture was partitioned between 100ml of methylene chloride and 100ml of 25% saturated aqueous sodium bicarbonate solution and the layers were separated. The aqueous layer was washed with 2 x 100ml of methylene chloride and acidified to pH 2 with 2N aqueous hydrochloric acid solution. The resulting mixture was extracted with 2 x 100ml of ethyl acetate; the extracts were combined, washed with 50ml of saturated aqueous sodium chloride solution, dried over magnesium sulfate, and concentrated *in vacuo* to afford 2.30g (77%) of the title compound as an oil that was used in the following step without further purification. IR Spectrum (neat,  $\text{cm}^{-1}$ ): 2111, 1724.  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ )  $\delta$  5.06 (1H, s), 7.08-7.45 (4H, m), 8.75 (1H, br s).

Step D: (S)-(4-Fluorophenyl)glycine

A mixture of 2.30g (11.8mmol) of (S)-azido-(4-fluorophenyl)acetic acid (from Step C), 2.50mg 10% palladium on carbon catalyst and 160ml 3:1 v/v water/acetic acid was stirred under an atmosphere of hydrogen for 5 18 hours. The reaction mixture was filtered through Celite and the flask and filter cake were rinsed well with about 1 litre of 3:1 v/v water/acetic acid. The filtrate was concentrated *in vacuo* to about 50ml of volume. 300ml of toluene was added and the mixture concentrated to afford a solid. The solid was suspended in 1:1 v/v methanol/ether, filtered and dried to 10 afford 1.99g (100%) of the title compound. <sup>1</sup>H NMR (400MHz, D<sub>2</sub>O+ NaOD) δ 3.97 (1H, s), 6.77 (2H, app t, J=8.8Hz), 7.01 (2H, app t, J=5.6Hz).

Via Resolution:

Step A' (4-Fluorophenyl)acetyl chloride

15 A solution of 150g (0.974mol) of 4-(fluorophenyl)acetic acid and 1ml of N,N-dimethylformamide in 500ml of toluene at 40°C was treated with 20ml of thionyl chloride and heated to 40°C. An additional 61.2ml of thionyl chloride was added dropwise over 1.5 hours. After the addition, the solution was heated at 50°C for 1 hour, the solvent was removed *in* 20 *vacuo* and the residual oil was distilled at reduced pressure (1.5mmHg) to afford 150.4g (89.5%) of the title compound, bp = 68-70°C.

Step B': Methyl 2-bromo-3-(4-fluorophenyl)acetate

25 A mixture of 150.4g (0.872mol) of 4-(fluorophenyl)acetyl chloride (from Step A') and 174.5g (1.09mol) of bromine was irradiated at 40-50°C with a quartz lamp for 5 hours. The reaction mixture was added dropwise to 400ml of methanol and the solution was stirred for 16 hours. The solvent was removed *in vacuo* and the residual oil was distilled at reduced pressure (1.5mmHg) to afford 198.5g (92%) of the title compound, bp = 30 106-110°C.

**Step C': Methyl (±)-(4-fluorophenyl)glycine**

A solution of 24.7g (0.1mol) of methyl 2-bromo-2-(4-fluorophenyl)acetate (from Step B') and 2.28g (0.01mol) of benzyl triethylammonium chloride in 25ml of methanol was treated with 6.8g (0.105mol) of sodium azide and the resulting mixture was stirred for 20 hours at room temperature. The reaction mixture was filtered; the filtrate was diluted with 50ml of methanol and hydrogenated in the presence of 0.5g of 10% Pd/C at 50 psi for 1 hour. The solution was filtered and the solvent removed *in vacuo*. The residue was partitioned between 10% aqueous sodium carbonate solution and ethyl acetate. The organic phase was washed with water, saturated aqueous sodium chloride solution dried over magnesium sulfate and concentrated *in vacuo* to afford 9.8g of the title compound as an oil.

**Step D': Methyl (S)-(4-fluorophenyl)glycinate**

A solution of 58.4g of methyl (±) 4-(fluorophenyl)glycinate (from Step C') in 110ml of 7:1 v/v ethanol/water was mixed with a solution of 28.6g (0.0799mol) of O,O'-(+)-dibenzoyltartaric acid ((+)-DBT) (28.6g, 0.0799mol) in 110ml of 7:1 v/v ethanol:water and the resulting solution was allowed to age at room temperature. Ethyl acetate (220ml) was added after crystallisation was complete and the resulting mixture was cooled to -20°C and filtered to afford 32.4g of methyl (S)-(4-fluorophenyl)glycinate, (+)-DBT salt (ee = 93.2%). The mother liquors were concentrated *in vacuo* and the free base was liberated by partitioning between ethyl acetate and aqueous sodium carbonate solution. A solution of free base, so obtained, in 110ml of 7:1 v/v ethanol/water was mixed with a solution of 28.6g (0.0799mol) of O,O'-(-)-dibenzoyltartaric acid ((-)-DBT) (28.6g, 0.0799mol) in 110ml of 7:1 v/v ethanol:water and the resulting solution was allowed to age at room temperature. Ethyl acetate (220ml) was added after crystallisation was complete and the resulting mixture was cooled to -20°C and filtered to afford 47.0g of methyl (R)-(4-fluorophenyl)glycinate,

(-)-DBT salt (ee = 75.8%). Recycling of the mother liquors and addition of (+)-DBT gave a second crop of 7.4g of (S)-(4-fluorophenyl)glycinate, (+)-DBT salt (ee = 96.4%). The two crops of the (S)-amino ester (39.8g) were combined in 200ml of 7:1 v/v ethanol/water, heated for 30 minutes and cooled to room temperature. Addition of ethyl acetate, cooling, and filtration afforded 31.7g of (S)-(4-fluorophenyl)glycinate, (+)-DBT salt (ee > 98%). Enantiomeric excess was determined by chiral HPLC (Crownpak CR(+) 5% MeOH in aq HClO<sub>4</sub> pH2 1.5ml/min 40°C 200nm).

A mixture of 17.5g of (S)-(4-fluorophenyl)glycinate, (+)-DBT salt and 32ml of 5.5N HCl (32ml) was heated at reflux for 1.5 hours. The reaction mixture was concentrated *in vacuo* and the residue was dissolved in 40ml of water. The aqueous solution was washed (3 x 30ml of ethyl acetate) and the layers were separated. The pH of the aqueous layer was adjusted to 7 using ammonium hydroxide and the precipitated solid was filtered to afford 7.4g of the title compound (ee = 98.8%).

## DESCRIPTION 2

### 4-Benzyl-3-(S)-(4-fluorophenyl)-2-morpholinone

#### 20 Step A: N-Benzyl-(S)-(4-fluorophenyl)glycine

A solution of 1.87g (11.05mmol) of (S)-(4-fluorophenyl)-glycine (from Description 1) and 1.12ml (11.1mmol) of benzaldehyde in 11.1ml of 1N aqueous sodium hydroxide solution and 11ml of methanol at 0°C was treated with 165mg (4.4mmol) of sodium borohydride. The cooling bath was removed and the resulting mixture was stirred at room temperature for 30 minutes. Second portions of benzaldehyde (1.12ml (11.1mmol)) and sodium borohydride (165mg (4.4mmol)) were added to the reaction mixture and stirring was continued for 1.5 hours. The reaction mixture was partitioned between 100ml of ether and 50ml of water and the layers were separated. The aqueous layer was separated and filtered to remove a small amount of insoluble material. The filtrate was acidified to pH 5

with 2N aqueous hydrochloric acid solution and the solid that had precipitated was filtered, rinsed well with water, then ether, and dried to afford 1.95g of the title compound. <sup>1</sup>H NMR (400MHz, D<sub>2</sub>O + NaOD) δ 3.33 (2H, AB q, J=8.4Hz), 3.85 (1H, s), 6.79-7.16 (4H, m).

5

**Step B: 4-Benzyl-3-(S)-(4-fluorophenyl)-2-morpholinone**

A mixture of 1.95g (7.5mmol) of N-benzyl (S)-(4-fluorophenyl)glycine, 3.90ml (22.5mmol) of N,N-diisopropyl-ethylamine, 6.50ml (75.0mmol) of 1,2-dibromoethane and 40ml of N,N-dimethylformamide was stirred at 100°C for 20 hours (dissolution of all solids occurred on warming). The reaction mixture was cooled and concentrated *in vacuo*. The residue was partitioned between 250ml of ether and 100ml of 0.5N potassium hydrogen sulfate solution and the layers were separated. The organic layer was washed with 100ml of saturated aqueous sodium bicarbonate solution, 3 x 150ml of water, dried over magnesium sulfate, and concentrated *in vacuo*. Flash chromatography on 125g of silica gel using 3:1 v/v hexanes/ether as the eluant afforded 1.58g (74%) of the title compound as an oil. <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>) δ 2.65 (1H, dt, J=3.2, 12.8Hz), 3.00 (1H, dt, J=12.8, 2.8Hz), 3.16 (1H, d, J=13.6Hz), 3.76 (1H, d, J=13.6Hz), 4.24 (1H, s), 4.37 (1H, dt, J=13.2, 3.2Hz), 4.54 (1H, dt, J=2.8, 13.2Hz), 7.07-7.56 (9H, m).

10

15

20

**DESCRIPTION 3**

**4-Benzyl-2-(R)-(3,5-bis(trifluoromethyl)benzoyloxy)-3-(S)-(4-fluorophenyl)morpholine**

25

A solution of 2.67g (10.0mmol) of 4-benzyl-3-(S)-(4-fluorophenyl)-2-morpholinone (Description 2) in 40ml of dry THF was cooled to -78°C. The cold solution was treated with 12.5ml of 1.0M L-Selectride® solution in THF, maintaining the internal reaction temperature below -70°C. The resulting solution was stirred cold for 45 minutes and the reaction was charged with 3.60ml(20.0mmol) of 3,5-bis(trifluoromethyl)benzoyl

30

- chloride. The resulting yellow mixture was stirred cold for 30 minutes and the reaction was quenched with 50ml of saturated aqueous sodium bicarbonate solution. The quenched mixture was partitioned between 300ml of ether and 50ml of water and the layers were separated. The
- 5 organic layer was dried over magnesium sulfate. The aqueous layer was extracted with 300ml of ether; the extract was dried and combined with the original organic layer. The combined organics were concentrated *in vacuo*. Flash chromatography on 150g of silica gel using 37:3 v/v hexanes/ether as the eluant afforded 4.06g (80%) of the title compound as a solid.
- 10 <sup>1</sup>H NMR (200MHz, CDCl<sub>3</sub>) δ 2.50 (1H, dt, J=3.4, 12.0Hz), 2.97 (1H, app d, J=12.0Hz), 2.99 (1H, d, J=13.6Hz), 3.72-3.79 (1H, m), 3.82 (1H, d, J=2.6Hz), 4.00 (1H, d, J=13.6Hz), 4.20 (dt, J=2.4, 11.6Hz), 6.22 (1H, d, J=2.6Hz), 7.22-7.37 (7H, m), 7.57 (2H, app d, J=6.8Hz), 8.07 (1H, s), 8.47 (2H, s). MS (FAB) m/z 528 (M+H, 25%), 270 (100%).
- 15 Analysis Calcd. for C<sub>28</sub>H<sub>20</sub>F<sub>7</sub>NO<sub>3</sub>: C, 59.21; H, 3.82; N, 2.66; F, 25.21; Found: C, 59.06; H, 4.05; N, 2.50; F, 25.18%.

#### DESCRIPTION 4

- 20 4-Benzyl-2-(R)-(1-(3,5-bis(trifluoromethyl)phenyl)ethenyloxy)-3-(S)-(4-fluorophenyl)morpholine

##### Step A: Dimethyl titanocene

- A solution of 2.49g (10.0mmol) of titanocene dichloride in 50ml of ether in the dark at 0°C was treated with 17.5ml of 1.4M methyllithium
- 25 solution in ether maintaining the internal temperature below 5°C. The resulting yellow/orange mixture was stirred at room temperature for 30 minutes and the reaction was quenched by slowly adding 25g of ice. The quenched reaction mixture was diluted with 50ml of ether and 25ml of water and the layers were separated. The organic layer was dried over
- 30 magnesium sulfate and concentrated *in vacuo* to afford 2.03g (98%) of the title compound as a light-sensitive solid. The dimethyl titanocene could be

stored as a solution in toluene at 0°C for at least 2 weeks without apparent chemical degradation. <sup>1</sup>H NMR (200MHz, CDCl<sub>3</sub>) δ -0.15 (6H, s), 6.06 (10H, s).

5    Step B: 4-Benzyl-2-(R)-(1-(3,5-bis(trifluoromethyl)phenyl)ethenyloxy)-3-(S)-(4-fluorophenyl)morpholine

A solution of the compound of Description 3 (2.50g, 4.9mmol) and 2.50g (12.0mmol) of dimethyl titanocene (from Step A) in 35ml of 1 litre v/v THF/toluene was stirred in an oil bath at 80°C for 16 hours. The  
10    reaction mixture was cooled and concentrated *in vacuo*. Flash chromatography on 150g of silica gel using 3:1 v/v hexanes/methylene chloride as the eluant afforded 1.71g (69%) of the title compound as a solid. An analytical sample was obtained via recrystallisation from isopropanol: <sup>1</sup>H NMR (400MHz, CDCl<sub>3</sub>) δ 2.42 (1H, dt, J=3.6, 12.0Hz),  
15    2.90 (1H, app d, J=12.0Hz), 2.91 (1H, d, J=13.6Hz), 3.62-3.66 (1H, m), 3.72 (1H, d, J=2.6Hz), 3.94 (1H, d, J=13.6Hz), 4.09 (1H, dt, J=2.4, 12.0Hz), 4.75 (1H, d, J=3.2Hz), 4.82 (1H, d, J=3.2Hz), 5.32 (1H, d, J=2.6Hz), 7.09 (2H, t, J=8.8Hz), 7.24-7.33 (5H, m), 7.58-7.62 (2H, m), 7.80 (1H, s), 7.90 (2H, s); MS (FAB) 526 (M+H, 75%), 270 (100%).  
20    Analysis Calcd. for C<sub>27</sub>H<sub>22</sub>F<sub>7</sub>NO<sub>2</sub>: C, 61.72; H, 4.22; N, 2.67; F, 25.31; Found: C, 61.79; H, 4.10; N, 2.65; F, 25.27%.

DESCRIPTION 5

25    2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)morpholine

The compound of Description 4 (4.0g) was dissolved in ethyl acetate (50ml) and isopropanol (16ml). To this solution was added palladium on charcoal (1.5g) and the mixture was hydrogenated at 40 psi for 36h. The catalyst was removed by filtration through Celite and the solvents were  
30    removed *in vacuo*. The residue was purified by flash chromatography on silica using 100% ethyl acetate and then 1-10% methanol in ethyl acetate.

This afforded isomer A 500mg (15%) and isomer B 2.6g (80%) as clear oils - isomer B crystallised on standing. For the title compound:  $^1\text{H}$  NMR (400MHz,  $\text{CDCl}_3$ )  $\delta$  1.16 (3H, d,  $J=6.8\text{MHz}$ ), 1.80 (1H, br s), 3.13 (1H, dd,  $J=3.2, 12.4\text{Hz}$ ), 3.23 (1H, dt,  $J=3.6, 12.4\text{Hz}$ ), 3.63 (1H, dd,  $J=2.4, 11.2\text{Hz}$ ),  
5 4.01 (1H, d,  $J=2.4\text{Hz}$ ), 4.13 (1H, dt,  $J=3.2, 12.0\text{Hz}$ ), 4.42 (1H, d,  $J=2.4\text{Hz}$ ), 4.19 (1H, q,  $J=6.8\text{Hz}$ ), 7.04-7.09 (2H, m), 7.27-7.40 (4H, m), 7.73 (1H, s); MS (FAB) 438 ( $\text{M}+\text{H}$ , 75%), 180 (100%).

HCl salt formation. To a solution of the free base (0.77g) in diethyl ether (10ml) was added 1M-HCl in methanol (1.75ml). The solution was  
10 evaporated to dryness and on addition of diethyl ether crystals formed. The solution was filtered and the residue washed with diethyl ether to give the title compound hydrochloride salt mp 248-250°C.

Analysis Calcd. for  $\text{C}_{20}\text{H}_{18}\text{F}_7\text{NO}_2\cdot\text{HCl}$ : C, 50.70; H, 4.04; N, 2.96; Cl, 7.48;  
Found: C, 50.46; H, 3.85; N, 3.01; Cl, 7.31%.

15

#### DESCRIPTION 6

##### 4-Benzyl-3-(S)-(4-fluorophenyl)-2-(R)-(3-fluoro-5-(trifluoromethyl)benzoyloxy)morpholine

The title compound was prepared from the reaction of the compound  
20 of Description 2 with 3-fluoro-5-(trifluoromethyl)benzoyl chloride according to the procedure illustrated in Description 3.  $^1\text{H}$  NMR (360MHz,  $\text{CDCl}_3$ )  $\delta$  2.50 (1H, dt,  $J=3.3, 12.0\text{Hz}$ ), 2.96 (1H, d,  $J=12.0\text{Hz}$ ), 2.98 (1H, d,  $J=13.6\text{Hz}$ ), 3.75 (1H, dd,  $J=1.7, 11.5\text{Hz}$ ), 3.80 (1H, d,  $J=2.5\text{Hz}$ ), 3.92 (1H, d,  $J=13.6\text{Hz}$ ), 4.19 (1H, dt,  $J=2.1, 12.0\text{Hz}$ ), 6.20 (1H, d,  $J=2.5\text{Hz}$ ), 6.99 (2H, t,  $J=8.7\text{Hz}$ ), 7.2-7.37 (5H, m), 7.51-7.55 (3H, m), 7.89  
25 (1H, d,  $J=8.4\text{Hz}$ ), 8.09 (1H, s). MS ( $\text{CI}^+$ )  $m/z$  478 ( $\text{M}^++1$ , 100%).

Analysis Calcd. for  $\text{C}_{25}\text{H}_{20}\text{F}_5\text{NO}_3$ : C, 62.88; H, 4.23; N, 2.93;  
Found: C, 62.59; H, 4.03; N, 3.07%.

30



NMR (360MHz, CDCl<sub>3</sub>) δ 2.42 (1H, dt, J=3.6, 12.0Hz), 2.90 (1H, d, J=12.0Hz), 2.91 (1H, d, J=13.6Hz), 3.60-3.62 (1H, m), 3.72 (1H, d, J=2.6Hz), 3.92 (1H, d, J=13.6Hz), 4.09 (1H, dt, J=2.4, 12.0Hz), 4.67 (1H, d, J=2.9Hz), 4.76 (1H, d, J=2.9Hz), 5.28 (1H, d, J=2.6Hz), 7.07 (2H, t, J=8.7Hz), 7.2-7.37 (7H, m), 7.53 (1H, s), 7.57-7.61 (2H, m). MS (CI<sup>+</sup>) 476 (M+1, 100%).

#### DESCRIPTION 8

15 3-(S)-(4-Fluorophenyl)-2-(R)-(1-(R)-(3-fluoro-5-(trifluoromethyl)phenyl)ethoxy)morpholine

The compound of Description 7 was hydrogenated according to the method illustrated in Description 5. This afforded a mixture of 2 epimeric products isomer A and isomer B (the major product) as clear oils. For the title compound: <sup>1</sup>H NMR (360MHz, CDCl<sub>3</sub>) δ 1.42 (3H, d, J=6.6Hz), 1.91 (1H, s), 3.11 (1H, dd, J=3.2, 12.4Hz), 3.22 (1H, dt, J=3.6, 12.4Hz), 3.58-3.62 (1H, m), 4.01 (1H, d, J=2.3Hz), 4.11 (1H, dt, J=3.2, 12.0Hz), 4.41 (1H, d, J=2.3Hz), 4.80 (1H, q, J=6.6Hz), 6.41 (1H, d, J=9.2Hz), 6.86 (1H, s), 7.02 (2H, t, J=8.7Hz), 7.08 (2H, d, J=9.2Hz), 7.21-7.26 (2H, m). MS (CI<sup>+</sup>) m/z 387 (M+1, 100%).

25 Analysis Calcd. for C<sub>19</sub>H<sub>18</sub>F<sub>5</sub>NO<sub>2</sub>: C, 58.91; H, 4.69; N, 3.62;  
Found: C, 58.88; H, 4.81; N, 3.76%.

#### DESCRIPTION 9

30 2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy-4-(5-(dimethylaminomethyl)-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine

Method A

- 5 a) 2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy-3-(S)-(4-fluorophenyl)-4-propargylmorpholine

Propargyl bromide (1.9ml) was added to a stirred mixture of the compound of Description 5 (5g) and potassium carbonate (4.76g) in dry dimethylformamide at 23°C. After 15 min the reaction mixture was  
 10 diluted with water (250ml) and extracted with ethyl acetate (3 x 100ml). The combined organic phases were washed with brine (1 x 100ml) then dried (K<sub>2</sub>CO<sub>3</sub>) and concentrated to leave an oil. This was purified by chromatography on silica using ethyl acetate in hexane (1:9 then 1:4) as eluent to afford the title compound as an oil. <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ  
 15 1.50 (3H, d, J=6.6Hz), 2.21 (1H, s), 2.84 (1H, d, J=11.1Hz), 2.97 (1H, td, J=3.2, 11.7Hz), 3.26 (2H, d, J=1.8Hz), 3.62 (1H, d, J=2.2Hz), 3.71 (1H, dd, J=2.3, 11.1Hz), 4.33 (2H, m), 4.89 (1H, q, J=6.6Hz), 7.03 (2H, t, J=8.6Hz), 7.18 (2H, s), 7.38 (2H, br s), 7.63 (1H, s). MS (CI<sup>+</sup>) m/z 476 (MH<sup>+</sup>, 100%).

- 20 b) 2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-4-(4-dimethylamino-4-oxo-but-2-ynyl)-3-(S)-(4-fluorophenyl)morpholine

A mixture of N,N-dimethylcarbamoyl chloride (0.195ml), cuprous iodide (2mg), bis(triphenylphosphine)palladium (II) chloride (2mg), triphenylphosphine (3mg) and the compound described in (a) above (1g) in  
 25 triethylamine (4ml) was heated at 90°C for 5h in an inert atmosphere. The mixture was cooled to 23°C and methanol (1ml) was added and the solvent was removed *in vacuo*. The residue was partitioned between water and ethyl acetate and the layers were separated. The aqueous phase was extracted with ethyl acetate (2 x 20ml). The combined organic  
 30 phases were washed with water, brine, dried (MgSO<sub>4</sub>) and concentrated to leave an oil. The residue was purified by chromatography on silica using

ethyl acetate in hexane (1:1) then ethyl acetate as eluant to provide the title compound as an oil. <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ 1.49 (3H, d, J=6.6Hz), 2.84-3.06 (2H, m), 3.00 (3H, s), 3.17 (3H, s), 3.44 (2H, s), 3.64 (1H, br s), 3.73 (1H, dd, J=2.0, 11.1Hz), 4.33 (2H, m), 4.88 (1H, q, J=6.6Hz), 7.03 (2H, t, J=8.7Hz), 7.17 (2H, s), 7.38 (2H, br s), 7.63 (1H, s). MS (CI<sup>+</sup>) m/z 547 (MH, 100%).

c) 2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-4-(5-N,N-dimethylcarboxamido-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine

A mixture of the compound described in (b) above (1.1g) and sodium azide (0.65g) in dimethylsulphoxide (7.5ml) was heated at 70°C for 17h. The mixture was cooled to 23°C and excess dimethylsulphoxide was removed by distillation *in vacuo*. The residue was partitioned between brine and ethyl acetate. The layers were separated and the organic layer was washed with brine (2 x 20ml) then dried (MgSO<sub>4</sub>) and concentrated to leave an oil. This was purified by chromatography on silica using ethyl acetate in hexane (1:2 then 1:1) and then ethyl acetate as eluent to provide the title compound as a pale yellow foam. <sup>1</sup>H NMR (360MHz, CDCl<sub>3</sub>) δ 1.47 (3H, d, J=6.6Hz), 2.64 (1H, m), 2.90 (1H, d, J=11.6Hz), 3.09 (3H, s), 3.34 (3H, s), 3.65 (3H, m), 3.92 (1H, d, J=15.5Hz), 4.27 (1H, td, J=2.1, 9.5Hz), 4.35 (1H, d, J=2.6Hz), 4.89 (1H, q, J=6.6Hz), 7.01 (2H, t, J=8.7Hz), 7.16 (2H, s), 7.39 (2H, br s), 7.64 (1H, s). m/z 590 (MH, 100%).

d) 2-(R)-(1-(R)-3,5-Bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(dimethylaminomethyl)-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine

Lithium aluminium hydride (0.47ml, 1M in tetrahydrofuran) was added dropwise to a solution of the compound described in (c) above (0.11g) in dry tetrahydrofuran (1ml) under an inert atmosphere at 23°C. After 30 min sodium hydroxide (10 drops, 1M) was added followed by

water (5 drops). Ethyl acetate (50ml) was then added and the resulting mixture was filtered through a pad of Hyflo™. The filtrate was concentrated *in vacuo* and the residue was purified by chromatography on silica using ethyl acetate in methanol (9:1 then 4:1) as eluant to provide the title compound as a foam. <sup>1</sup>H NMR (360MHz, CDCl<sub>3</sub>) δ 1.44 (3H, d, J=6.6Hz), 2.25 (6H, s), 2.57 (1H, td, J=3.4, 8.55Hz), 2.90 (1H, d, J=11.7Hz), 3.25 (1H, d, J=14.0Hz), 3.43 (1H, d, J=13.6Hz), 3.45 (1H, d, J=2.2Hz), 3.53 (1H, d, J=13.6Hz), 3.61 (1H, d, J=11.2Hz), 3.78 (1H, d, J=14.0Hz), 4.22 (1H, t, J=9.3Hz), 4.32 (1H, d, J=2.2Hz), 4.86 (1H, q, J=6.6Hz), 7.06 (2H, t, J=8.7Hz), 7.16 (2H, s), 7.48 (2H, br s), 7.63 (1H, s). m/z 576 (MH).

#### Method B

2-(R)-1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(4-chlorobut-2-ynyl)morpholine

a) A solution of the product of Description 5 (free base, 5g) in N,N-dimethylformamide (20ml) was slowly added to a heated (50°C) solution of 1,4-dichlorobut-2-yne (2.2ml) and potassium carbonate (4.8g) in N,N-dimethylformamide (20ml). The solution was heated for a further 5h at 50°C and then the solvent removed *in vacuo*. To the residue was added water (400ml) and the product extracted into ethyl acetate (3 x 150ml). The combined organic phase washed with water, saturated brine and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue chromatographed on silica gel (eluting with 10% ethyl acetate in petroleum ether bp 60-80°C) to give the title compound. <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ 1.41 (3H, d, J=6.6Hz), 2.80 (1H, app. t, J=10.8Hz), 2.87 (1H, td, J=3.5Hz, 11.7Hz), 3.22 (2H, t, J=1.9Hz), 3.52 (1H, d, J=2.8Hz), 3.68 (1H, d, J=1.4Hz, 11.1Hz), 4.00 (2H, t, J=1.9Hz), 4.22-4.32 (2H, m), 4.81 (1H, q, J=6.6Hz), 6.96 (2H, t, J=8.7Hz), 7.10 (2H, s), 7.31 (2H, br s), 7.56 (1H, s). m/z (CI<sup>+</sup>) 524 (M+H, 100%).

b) N-(4-Azidobut-2-ynyl)-2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)morpholine

5 To a solution of 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(4-chlorobut-2-ynyl)morpholine (4g) in dimethyl sulphoxide (17ml) was added sodium azide (0.562g). The solution was stirred for 20h and aqueous ammonium chloride and ethyl acetate were added. The organic phase was washed with water (2 times),  
10 saturated brine and dried (MgSO<sub>4</sub>). The solvent was removed *in vacuo* and the residue chromatographed on silica gel (eluting with 20% ethyl acetate in petroleum ether bp 60-80°C) to give the title compound. <sup>1</sup>H NMR (360MHz, CDCl<sub>3</sub>) δ 1.48 (3H, s, J=6.6Hz), 2.87 (1H, app t, J=10.2Hz), 2.98 (1H, td, J=3.6, 11.7Hz), 3.35 (2H, t, J=1.9Hz), 3.61 (1H, d, J=2.8Hz), 3.72 (1H, dq, J=1.4Hz, 10.0Hz), 3.92 (2H, t, J=1.9Hz), 4.30-4.40  
15 (2H, m), 4.89(1H, q, J=6.6Hz), 7.03 (2H, t, J=8.7Hz), 7.17 (2H, s), 7.27 (2H, br s), 7.63 (1H, s).

c) 2-(R)-(1-(R)-3,5-Bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(dimethylaminomethyl)-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine

20 Dimethylamine (approximately 10ml) was condensed at -80°C in a pressure tube and to this was added a solution of N-(4-azidobut-2-ynyl)-2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)morpholine (3.2g) in dioxan (15ml). The tube was sealed and the solution was heated at 90°C for 16h. The solution was evaporated to dryness and the residue chromatographed on silica gel (eluting with 5% methanol in dichloromethane containing 0.25% ammonia (SG. 0.88)) and the fractions containing the desired product were evaporated *in vacuo* to  
25 give the title compound. To a solution of this residue in diethyl ether was added 1M-HCl in methanol. The solution was evaporated to dryness and  
30

redissolved in diethyl ether to give crystals of the title compound hydrochloride salt m.p. 194-198°C,  $[\alpha]_D^{25} + 65.0^\circ$  (c=0.5, H<sub>2</sub>O). The crystals were found to be stable for at least five days at 40°C; at 40°C/75% relative humidity; at 80°C; and at 2000LUX.

5

#### DESCRIPTION 10

3-(S)-(4-Fluorophenyl)-2-(R)-(1-(R)-(3-fluoro-5-(trifluoromethyl)phenyl)ethoxy)-4-(5-pyrrolidinomethyl-1,2,3-triazol-4-yl)methylmorpholine

The title compound was prepared from the compound of Description 8 according to the method of Description 9, Method B, via the  
10 corresponding N-(4-azidobut-2-ynyl)morpholine and pyrrolidine. <sup>1</sup>H NMR (360MHz, CDCl<sub>3</sub>) δ 1.40 (3H, d, J=6.6Hz), 1.81 (4H, br s), 2.53-2.61 (5H, m), 2.89 (1H, d, J=11.7Hz), 3.27 (1H, d, J=14.0Hz), 3.45 (1H, d, J=2.8Hz), 2.59-3.63 (1H, m), 3.63 (1H, d, J=13.7Hz), 3.73 (1H, d, J=13.7Hz), 3.83  
15 (1H, d, J=14.0Hz), 4.21 (1H, dt, J=11.6, 2.1Hz), 4.32 (1H, d, J=2.8Hz), 4.76 (1H, q, J=6.5Hz), 6.37 (1H, d, J=9.1Hz), 6.80 (1H, s), 7.05-7.10 (3H, m), 7.46 (2H, br s). MS (CI<sup>+</sup>) 552 (M+1, 100%).

#### DESCRIPTION 11

20 3-(S)-(4-Fluorophenyl)-2-(R)-(1-(R)-(3-methylthio-5-(trifluoromethyl)phenyl)ethoxy)-4-(5-pyrrolidinomethyl-1,2,3-triazol-4-yl)methylmorpholine

The compound of Description 10 (0.5mmol) was heated to 120°C with sodium thiomethoxide (2.5mmol) in anhydrous DMF (10ml) for  
25 between 2-5 hours. The cooled solution was diluted with water (150ml), extracted with ethyl acetate (4 x 40ml), dried (MgSO<sub>4</sub>) and concentrated *in vacuo* to a crude oil which was purified by flash silica gel chromatography in 5-10% methanol/dichloromethane to yield the title compound as a foam (620mg, 81%). <sup>1</sup>H NMR (360MHz, CDCl<sub>3</sub>) δ 1.40 (3H,  
30 d, J=6.6Hz), 1.79 (4H, br s), 2.36 (3H, s), 2.5-2.6 (5H, m), 2.87 (1H, d, J=11.7Hz), 3.23 (1H, d, J=13.9Hz), 3.43 (1H, d, J=2.8Hz), 3.57-3.64 (2H,

m), 3.71 (1H, d, J=13.7Hz), 3.78 (1H, d, J=14.0Hz), 4.21 (1H, m), 4.33 (1H, d, J=2.8Hz), 4.74 (1H, q, J=6.5Hz), 6.71 (2H, s), 7.06 (2H, t, J=8.7Hz), 7.19 (1H, s), 7.47 (2H, br s); MS (ES<sup>+</sup>) m/z 580 (M+1, 100%).

5

# EXAMPLE 1

2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(5-(N,N-dimethylamino(N-oxide)methyl)-1,2,3-triazol-yl)methylmorpholine

To a heated (60°C) solution of 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(5-(N,N-dimethylaminomethyl)-1,2,3-triazol-4-yl)methylmorpholine (Description 9) (0.245g, 0.442mmol) dissolved in soya bean oil (6ml) was added a 30% solution of aqueous hydrogen peroxide (0.2ml) in portions over 4h. The solution was then cooled to room temperature for 16 hours and the resulting suspension applied to a column containing silica and the products eluted with a gradient between dichloromethane and a mixture of dichloromethane:methanol:aqueous ammonia solution (200:50:1). This crude product was recrystallised from aqueous ethanol to give the title compound mp 139-140°C. m/z (CI<sup>+</sup>) 592 (M+H).

20 Analysis Calcd. for C<sub>26</sub>H<sub>28</sub>F<sub>7</sub>N<sub>5</sub>O<sub>3</sub> · 0.8H<sub>2</sub>O: C, 51.54; H, 4.92; N, 11.56;  
Found: C, 51.52; H, 4.71; N, 11.21%  
<sup>1</sup>H NMR (360MHz, CH<sub>3</sub>OH-d<sub>4</sub>) δ 1.47 (3H, d, J=6.58Hz), 2.54 (1H, td), 2.78 (1H, d, J=11.95Hz), 3.10 (3H, s), 3.15 (3H, s), 3.35 (1H, AB d, J=14.49Hz), 3.50 (1H, d), 3.63 (1H, br d, J=11.53Hz), 3.82 (1H, AB d, J=14.5Hz), 4.23 (1H, td), 4.29 (1H, AB d, J=13.55Hz), 4.36 (1H, d, J=2.69Hz), 4.49 (1H, AB d, J=13.43Hz), 4.96 (1H, q, J=6.48Hz), 7.08 (2H, t, J=8.7Hz), 7.35 (2H, s), 7.56 (2H, br t), 7.71 (1H, s).

2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-4-(5-(N,N-dimethylaminomethyl)-1,2,3-triazol-4-yl) methylmorpholine 1-oxide

Analysis Calcd. for  $C_{26}H_{28}F_7N_5O_3 \cdot 1.3H_2O$ : C, 50.78; H, 5.02; N, 11.39;  
Found: C, 50.67; H, 5.05; N, 11.15%

25 EXAMPLE 3

2-(R)-(1-(R)-(3,5-Bis(trifluoromethyl)phenyl)ethoxy)-3-(S)-(4-fluorophenyl)-  
4-(5-(N,N-dimethylamino(N-oxide)methyl)-1,2,3-triazol-4-yl)  
methylmorpholine N-oxide

From the chromatographic separation described in Example 1 was  
30 isolated the title compound  $m/z$  (CI<sup>+</sup>) 608 (M+H).



**3-(S)-(4-Fluorophenyl)-2-(R)-(1-(R)-(3-methylthio-5-(trifluoromethyl)phenyl)ethoxy)-4-(5-pyrrolidinomethyl-1,2,3-triazol-4-yl)methylmorpholine N-oxide**

Analysis Calcd. for $C_{22}H_{33}F_4N_5O_3S \cdot 2H_2O$ :	C, 53.24; H, 5.90; N, 11.09;
Found:	C, 53.24; H, 5.74; N, 10.76%

25


3-(S)-(4-Fluorophenyl)-2-(R)-(1-(R)-(3-methylthio-5-(trifluoromethyl)phenyl)ethoxy)-4-(5-pyrrolidino(N-oxide)methyl-1,2,3-triazol-4-yl)methylmorpholine N-oxide

From the chromatography described in Example 4. <sup>1</sup>H NMR (360MHz, CDCl<sub>3</sub>) δ 1.45 (3H, d, J=6.6Hz), 1.90-2.40 (4H, br m), 2.28 (3H, s), 3.24 (1H, d, J=12.9Hz), 3.35 (1H, m), 3.56-3.85 (5H, br m), 4.08 (1H, d,

J=13.0Hz), 4.50 (1H, d, J=13.1Hz), 4.56-4.71 (3H, m), 4.83 (1H, d, J=13.1Hz), 5.04 (1H, d, J=3.4Hz), 5.13 (1H, d, J=13.0Hz), 6.89 (2H, s), 6.99 (2H, br t), 7.16 (1H, s), 7.6-8.2 (~2H, vbr s); MS (ES<sup>+</sup>) 612 (M+1, 100%).

**CLAIMS:**

1. A compound of the formula (I):



(I)

wherein

R<sup>1</sup> is hydrogen, halogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, CF<sub>3</sub>, OCF<sub>3</sub>, NO<sub>2</sub>, CN, SR<sup>a</sup>, SOR<sup>a</sup>, SO<sub>2</sub>R<sup>a</sup>, CO<sub>2</sub>R<sup>a</sup>, CONR<sup>a</sup>R<sup>b</sup>, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl or C<sub>1-4</sub>alkyl substituted by C<sub>1-4</sub>alkoxy, wherein R<sup>a</sup> and R<sup>b</sup> each independently represent hydrogen or C<sub>1-4</sub>alkyl;

R<sup>2</sup> is hydrogen, halogen, C<sub>1-6</sub>alkyl, CF<sub>3</sub>, OCF<sub>3</sub> or C<sub>1-6</sub>alkoxy substituted by C<sub>1-4</sub>alkoxy;

R<sup>3</sup> is hydrogen, halogen, CF<sub>3</sub> or OCF<sub>3</sub>;

R<sup>4</sup> is hydrogen, halogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy, CF<sub>3</sub>, NO<sub>2</sub>, CN, SR<sup>a</sup>, SOR<sup>a</sup>, SO<sub>2</sub>R<sup>a</sup>, CO<sub>2</sub>R<sup>a</sup>, CONR<sup>a</sup>R<sup>b</sup>, C<sub>2-6</sub>alkenyl, C<sub>2-6</sub>alkynyl or C<sub>1-4</sub>alkyl substituted by C<sub>1-4</sub>alkoxy, wherein R<sup>a</sup> and R<sup>b</sup> are as previously defined;

R<sup>5</sup> is hydrogen, halogen, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkoxy substituted by C<sub>1-4</sub>alkoxy or CF<sub>3</sub>;

R<sup>6</sup> is hydrogen, C<sub>1-4</sub>alkyl, C<sub>3-7</sub>cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-4</sub>alkyl, or C<sub>2-4</sub>alkyl substituted by C<sub>1-4</sub>alkoxy or hydroxy;

R<sup>7</sup> is hydrogen, C<sub>1-4</sub>alkyl, C<sub>3-7</sub>cycloalkyl, C<sub>3-7</sub>cycloalkylC<sub>1-4</sub>alkyl, or C<sub>2-4</sub>alkyl substituted by one or two substituents selected from C<sub>1-4</sub>alkoxy, hydroxy or a 4, 5 or 6 membered heteroaliphatic ring containing one or two heteroatoms selected from N, O and S;

- 5 or R<sup>6</sup> and R<sup>7</sup>, together with the nitrogen atom to which they are attached, form a saturated or partially saturated heterocyclic ring of 4 to 7 ring atoms, which ring may optionally contain in the ring one oxygen or sulphur atom or a group selected from NR<sup>8</sup>, S(O) or S(O)<sub>2</sub> and which ring may be optionally substituted by one or two groups selected from hydroxy, 10 C<sub>1-4</sub>alkyl, hydroxyC<sub>1-4</sub>alkyl, C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl, oxo, COR<sup>a</sup> or CO<sub>2</sub>R<sup>a</sup> where R<sup>a</sup> is as previously defined;

or R<sup>6</sup> and R<sup>7</sup> together with the nitrogen atom to which they are attached, form a non-aromatic azabicyclic ring system of 6 to 12 ring atoms;

- 15 or Z, R<sup>6</sup> and the nitrogen atom to which they are attached form a heteroaliphatic ring of 4 to 7 ring atoms which may optionally contain an oxygen ring atom;

R<sup>8</sup> is hydrogen, C<sub>1-4</sub>alkyl, hydroxyC<sub>1-4</sub>alkyl or C<sub>1-4</sub>alkoxyC<sub>1-4</sub>alkyl;

- R<sup>9a</sup> and R<sup>9b</sup> are each independently hydrogen or C<sub>1-4</sub>alkyl, or R<sup>9a</sup> and 20 R<sup>9b</sup> are joined so, together with the carbon atoms to which they are attached, there is formed a C<sub>5-7</sub> ring;

Het is a 5- or 6-membered heterocyclic ring containing 2 or 3 nitrogen atoms optionally substituted by =O, =S or a C<sub>1-4</sub>alkyl group;

- X is an alkylene chain of 1 to 4 carbon atoms optionally substituted 25 by oxo;

Y is a C<sub>1-4</sub>alkyl group optionally substituted by hydroxy;

Z is C<sub>1-6</sub>alkylene or C<sub>3-7</sub>cycloalkylene;

m is 0 or 1; and

n is 0 or 1, where the sum total of n+m is 1 or 2;

- 30 or a pharmaceutically acceptable salt thereof.

2. A compound as claimed in claim 1 wherein  $R^1$  is hydrogen,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy, halogen or  $CF_3$ .


3. A compound as claimed in claim 1 or claim 2 wherein  $R^2$  is  
5 hydrogen,  $C_{1-4}$ alkyl,  $C_{1-4}$ alkoxy, halogen or  $CF_3$ .

4. A compound as claimed in any one of claims 1 to 3 wherein  $R^3$  is hydrogen, fluorine, chlorine or  $CF_3$ .

10 5. A compound as claimed in any one of claims 1 to 4 wherein m is 1.

6. A compound of the formula (Ia) or a pharmaceutically acceptable salt thereof:

15



(Ia)

wherein  $A^1$  is fluorine or  $CF_3$ ;

$A^2$  is fluorine or  $CF_3$ ;

20  $A^3$  is fluorine or hydrogen;


and  $R^6$ ,  $R^7$ ,  $X$ ,  $Y$ ,  $Z$ ,  $Het$ ,  $m$  and  $n$  are as defined in claim 1.

7. A compound as claimed in any one of claims 1 to 6 wherein Y is a C<sub>1-4</sub>alkyl group.

5 8. A compound as claimed in any one of claims 1 to 7 wherein X is CH<sub>2</sub>, CH(CH<sub>3</sub>) or CH<sub>2</sub>CH<sub>2</sub>.


9. A compound as claimed in any one of claims 1 to 8 wherein the group Het-ZN(O)<sub>m</sub>R<sup>6</sup>R<sup>7</sup> is selected from:

10



10. A compound as claimed in claim 9 wherein the group Het-ZN(O)<sub>m</sub>R<sup>6</sup>R<sup>7</sup> is:

15



11. A compound a claimed in any one of claims 1 to 10 wherein Z is CH<sub>2</sub> or CH<sub>2</sub>CH<sub>2</sub> and NR<sup>6</sup>R<sup>7</sup> is amino, methylamino, dimethylamino, diethylamino, azetidiny, pyrrolidino and morpholino.

12. A compound selected from:

- 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(N,N-dimethylamino(N-oxide)methyl)-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine;
- 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(N,N-dimethylaminomethyl)-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine N-oxide;
- 2-(R)-(1-(R)-(3,5-bis(trifluoromethyl)phenyl)ethoxy)-4-(5-(N,N-dimethylamino(N-oxide)methyl)-1,2,3-triazol-4-yl)methyl-3-(S)-(4-fluorophenyl)morpholine N-oxide;
- 3-(S)-(4-fluorophenyl)-2-(R)-(1-(R)-(3-methylthio-5-(trifluoromethyl)phenyl)ethoxy)-4-(5-pyrrolidinomethyl-1,2,3-triazol-4-yl)methylmorpholine N-oxide;
- 3-(S)-(4-fluorophenyl)-2-(R)-(1-(R)-(3-methylthio-5-(trifluoromethyl)phenyl)ethoxy)-4-(5-pyrrolidino(N-oxide)methyl-1,2,3-triazol-4-yl)methylmorpholine N-oxide;
- or a pharmaceutically acceptable salt thereof.

12. A compound as claimed in any preceding claim for use in therapy.

20


13. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 11 in association with a pharmaceutically acceptable carrier or excipient.

14. The use of a compound as claimed in any one of claims 1 to 11 for the manufacture of a medicament for the treatment or prevention of a physiological disorder associated with an excess of tachykinins.

15. The use of a compound as claimed in any one of claims 1 to 11 for the manufacture of a medicament for the treatment or prevention of pain, inflammation, migraine, emesis or postherpetic neuralgia.

16. A process for the preparation of a compound as claimed in claim 1 which comprises:  
oxidation of one or both of the nitrogen atoms drawn in formula (II):

5



(II)

said process being followed, where necessary, by the removal of any protecting group where present;

- and when the compound of formula (I) is obtained as a mixture of enantiomers or diastereoisomers, optionally resolving the mixture to obtain the desired enantiomer;

10

and/or, if desired, converting the resulting compound of formula (I) or a salt thereof, into a pharmaceutically acceptable salt thereof.





The  
Patent  
Office

62

Application No: GB 9613003.4  
Claims searched: 1-16

Examiner: Roy Honeywood  
Date of search: 5 September 1996

**Patents Act 1977**  
**Search Report under Section 17**

**Databases searched:**

UK Patent Office collections, including GB, EP, WO & US patent specifications, in:  
UK CI (Ed.O): C2C  
Int CI (Ed.6): C07D  
Other: ONLINE: CAS

**Documents considered to be relevant:**

Category	Identity of document and relevant passage	Relevant to claims
	None	

X	Document indicating lack of novelty or inventive step	A	Document indicating technological background and/or state of the art.
Y	Document indicating lack of inventive step if combined with one or more other documents of same category.	P	Document published on or after the declared priority date but before the filing date of this invention.
&	Member of the same patent family	E	Patent document published on or after, but with priority date earlier than, the filing date of this application.

